

Exact results for sheared polar active suspensions with variable liquid crystalline order

Cite as: J. Chem. Phys. **150**, 104902 (2019); <https://doi.org/10.1063/1.5080343>

Submitted: 07 November 2018 . Accepted: 18 February 2019 . Published Online: 08 March 2019

Aurore Loisy , Anthony P. Thompson, Jens Eggers, and Tanniemola B. Liverpool 



[View Online](#)



[Export Citation](#)



[CrossMark](#)

Exact results for sheared polar active suspensions with variable liquid crystalline order

Cite as: J. Chem. Phys. 150, 104902 (2019); doi: 10.1063/1.5080343

Submitted: 7 November 2018 • Accepted: 18 February 2019 •

Published Online: 8 March 2019



View Online



Export Citation



CrossMark

Aurore Loisy,  Anthony P. Thompson, Jens Eggers, and Tanniemola B. Liverpool^{a)} 

AFFILIATIONS

School of Mathematics, University of Bristol, Bristol BS8 1TW, United Kingdom

Note: This article is part of the Special Topic “Chemical Physics of Active Matter” in J. Chem. Phys.

^{a)} Author to whom correspondence should be addressed: t.liverpool@bristol.ac.uk

ABSTRACT

We consider a confined sheared active polar liquid crystal with a uniform orientation and study the effect of variations in the magnitude of polarization. Restricting our analysis to one-dimensional geometries, we demonstrate that with asymmetric boundary conditions, this system is characterized, macroscopically, by a linear shear stress vs. shear strain relationship that does not pass through the origin: At a zero strain rate, the fluid sustains a non-zero stress. Analytic solutions for the polarization, density, and velocity fields are derived for asymptotically large or small systems and are shown by comparison with precise numerical solutions to be good approximations for finite-size systems.

Published under license by AIP Publishing. <https://doi.org/10.1063/1.5080343>

I. INTRODUCTION

Swarms of bacteria,¹⁻⁴ mixtures of cytoskeletal filaments and motor proteins,⁵⁻⁷ and self-propelled colloids^{8,9} are all examples of active suspensions¹⁰⁻¹³ consisting of anisotropic self-driven particles dispersed in a passive liquid. Due to the orientable nature of their constituents, active suspensions can exhibit a long-range orientational order and are often referred to as active liquid crystals (LCs).^{10-12,14} While active LCs can exist in ordered phases typical of liquid crystals,¹⁵ they fundamentally differ from their passive counterparts, because each active particle transduces free energy into systematic movement, maintaining the system out of equilibrium.

As active particles interact with each other and with their surrounding environment, they are able to collectively generate motion and mechanical stresses at scales much larger than their individual size, endowing active materials with unusual mechanical properties. An example is the reduction of the apparent viscosity of bacterial suspensions under shear.¹⁶⁻²¹ Remarkably, upon increasing activity, the apparent viscosity can decrease until a value of zero is achieved, giving rise to superfluid-like behaviour.^{20,22} In the last decade, rheological measurements¹⁶⁻²⁰ have shown qualitative agreement with earlier theoretical predictions²³⁻²⁸ for the macroscopic mechanical properties of active suspensions. Yet, a more thorough understanding of the underlying mechanisms driving these systems

requires a more quantitative comparison of theoretical models with experiments.²⁹ Such a comparison is becoming possible as more detailed information, such as transient rheological behaviour²⁰ and velocity profiles,²² becomes accessible experimentally.

Many active materials, whether biological or synthetic, involve head-tail asymmetric particles and can exist in a polar phase. For such systems, the broken symmetry variable is the polarization vector which represents the local coarse-grained orientation of the particles. When an active polar LC is sheared, distortions in the orientation of the polarization field induce active stresses which in turn generate an extra flow (needed to maintain the stress balance). This mechanism allows the apparent viscosity (defined as the macroscopic viscosity at the scale of the system, as would be measured by using a rheometer) of active LCs to vanish or even become negative.²⁹ While previous theoretical analyses²⁹⁻³¹ have focused on how gradients in the *orientation* can induce the active stresses that lead to unconventional mechanical behaviour, here our focus is on how variations in the *magnitude* of LC order affect the mechanical properties of active LCs.

In this paper, we show that a gradient in the magnitude of polarization of active LCs induces, even in the *absence* of variations in orientation, flows that give rise to anomalous mechanics. Specifically, we examine the effect of a varying polarization magnitude on a one-dimensional confined active polar LC subjected to shear. We derive the analytical relationship between the stress and

the macroscopic strain rate, which shows, in particular, that this system experiences a non-zero shear stress at a zero shear strain rate (and conversely, a non-zero strain rate is required to maintain a zero stress).

With the assumption of a uniform orientation, the nonlinear governing equations become decoupled, so it becomes possible to obtain explicit solutions. We obtain exact analytical expressions for the velocity and density fields as a function of the polarization field, the latter being shown to be well approximated by asymptotic solutions in the limit of large or small systems. This is interesting because hydrodynamic equations for active LCs are highly nonlinear; hence, analytical solutions beyond linearization approximations are rare even in the simplest geometries. As a result, studies of sheared active LCs tend to be numerical.²⁹⁻³⁵

II. MODEL

We consider an active LC with the possibility of polar orientational order. At the continuum scale, its dynamics are described by a set of long-wavelength, long-time scale equations forming the now well-accepted hydrodynamic theory of active matter.^{11,14} The relevant hydrodynamic variables are the polarization vector \mathbf{p} , the particle number density ρ (for simplicity ρ is normalized by its equilibrium value), and the momentum is $\rho_m \mathbf{u}$, where ρ_m is the fluid mass density and \mathbf{u} is the fluid velocity.

The passive contributions to the equations of motion are customarily described as those arising from the nonequilibrium analog of the free energy for a passive polar LC. This free energy is given by^{11,30}

$$F = \int_r (f_n + f_p) dr, \quad (1)$$

$$f_n = \frac{a_2}{2} |\mathbf{p}|^2 + \frac{a_4}{4} |\mathbf{p}|^4 + \frac{K}{2} |\nabla \mathbf{p}|^2 + \frac{C}{2} (\rho - 1)^2, \quad (2)$$

$$f_p = B_1 (\rho - 1) \nabla \cdot \mathbf{p} + B_2 |\mathbf{p}|^2 \nabla \cdot \mathbf{p} + B_3 |\mathbf{p}|^2 \mathbf{p} \cdot \nabla \rho. \quad (3)$$

The contribution f_n is the free energy density of a nematic LC. The first two terms control the isotropic-polar transition: They favor a polar phase ($|\mathbf{p}|^2 = -a_2/a_4$) when $a_2 < 0$ and an isotropic phase ($|\mathbf{p}|^2 = 0$) when $a_2 > 0$. The third term describes the energy cost of deformation (K is the analog of the Frank constant for passive LCs), and the last term penalizes density variations (C is the compression modulus). The contribution f_p contains additional terms that break the $\mathbf{p} \rightarrow -\mathbf{p}$ symmetry and are allowed in a polar fluid.³⁶

The polarization and density dynamics are governed by

$$D_t \mathbf{p} = -\beta_p \mathbf{p} \cdot \nabla \mathbf{p} + \lambda \mathbf{E} \cdot \mathbf{p} - \Omega \cdot \mathbf{p} - \Gamma_{pp} \mathbf{h} - \Gamma_{cp} \mathbf{g} \quad (4)$$

and

$$D_t \rho = \nabla \cdot [-\rho \beta_c \mathbf{p} + \Gamma_{cp} \mathbf{h} + \Gamma_{cc} \mathbf{g}], \quad (5)$$

where $D_t = \partial_t + \mathbf{u} \cdot \nabla$, $E_{ij} = (\partial_i u_j + \partial_j u_i)/2$, $\Omega_{ij} = (\partial_i u_j - \partial_j u_i)/2$, $\mathbf{h} = \delta F / \delta \mathbf{p}$, and $\mathbf{g} = \nabla(\delta F / \delta \rho)$. The flow is assumed incompressible ($\nabla \cdot \mathbf{u} = 0$), and the flow field satisfies

$$\rho_m (\partial_t + u_i \partial_i) u_j = \partial_i \sigma_{ij}. \quad (6)$$

The stress tensor is given by

$$\sigma_{ij} = 2\eta E_{ij} + \sigma_{ij}^r + \sigma_{ij}^a, \quad (7)$$

where the first term is the dissipative contribution (η is the fluid viscosity), σ_{ij}^r is the reversible contribution (as in passive LCs), and σ_{ij}^a is the active contribution. The reversible stress is given by

$$\sigma_{ij}^r = -\Pi \delta_{ij} + \frac{\lambda}{2} (p_i h_j + p_j h_i) + \frac{1}{2} (p_i h_j - p_j h_i), \quad (8)$$

where Π is the pressure. The active stress is

$$\sigma_{ij}^a = \alpha \rho p_i p_j + \beta_\sigma \rho (\partial_i p_j + \partial_j p_i), \quad (9)$$

where the lowest order term $\sim \alpha$ has nematic symmetry, while the higher order term $\sim \beta_\sigma$ is present only in systems with polar symmetry. In the above equations, $\beta_{c,p}$ and $\beta_\sigma \Gamma_{pp}$ have the dimension of a velocity, and the associated terms arise in polar systems from the self-advection of active elements along \mathbf{p} .

Our geometry is similar to that used in prior work^{29,30} and is depicted in Fig. 1: A two-dimensional thin film of active LC of thickness L is sheared between two parallel walls moving in opposite directions with velocity magnitude V . We allow gradients only in the direction normal to the walls. Due to incompressibility and wall impermeability, the flow must be parallel to the wall: $\mathbf{u} = (u(z), 0)$, and we use no-slip boundary conditions: $u(L) = -u(0) = V$. The fluid is therefore subjected to a macroscopic shear strain rate $\dot{\gamma} = \int_0^L \partial_z u dz = 2V/L$.

To pick out the effects of variations in the amount of LC order only, we further assume that the polarization field is parallel to the walls, and only its (signed) magnitude is allowed to vary: $\mathbf{p} = (p(z), 0)$. Previous work has focused on the role of varying orientation with fixed magnitude, here by contrast, we fix the orientation and focus on the role of varying magnitude of \mathbf{p} only. Our choice of orientation is consistent with the expected boundary conditions on the walls. An attractive feature of this approximation is that it makes the problem tractable analytically due to the decoupling of the equations. Such an aligned state is physically relevant to situations where strong parallel anchoring is prescribed at the walls, provided that the coupling between the polarization orientation and the local shear is negligible,




FIG. 1. A thin film of active polar LC sheared between two moving no-slip walls. The polarization field is uniformly aligned with the walls, and only its magnitude p is allowed to vary.

that is, for systems which satisfy $\Gamma_{pp}K \gg UL$ (with U being a characteristic velocity scale for the flow). The governing equations reduce to

$$\partial_t p = -\Gamma_{pp}(a_2 + a_4 p^2)p + \Gamma_{pp}K\partial_z^2 p, \quad (10)$$

$$\partial_t \rho = \partial_z[-2b_2 p \partial_z p + (d + b_3 p^2)\partial_z \rho], \quad (11)$$

$$\rho_m \partial_t u = \partial_z \sigma, \quad (12)$$

with

$$\sigma = \eta \partial_z u + (\beta_\sigma \rho + mb_2 p^2)\partial_z p + \frac{m}{2}(b_1 - b_3 p^2)p \partial_z \rho, \quad (13)$$

where σ_{zx} is now simply denoted σ and we have introduced $d = \Gamma_{cc}C - \Gamma_{cp}B_1$, $b_{1,2,3} = \Gamma_{cp}B_{1,2,3}$, and $m = (1 - \lambda)/\Gamma_{cp}$. At the boundaries, the flux of ρ across the walls must be zero, and we require that the polarization vectors at the walls are antiparallel: $p(L) = -p(0) = p_{eq} = \sqrt{-a_2/a_4}$ (here, we assume $a_2 < 0$).

It is interesting to note that the coupling terms in the governing equations are those which break the $\mathbf{p} \rightarrow -\mathbf{p}$ symmetry. For a nematic fluid ($b_{1,2,3} = 0$, $\beta_\sigma = 0$), the equation for ρ reduces to a diffusion equation, and the expression of the stress only contains the viscous term. Therefore, in the simple configuration we consider here, a nematic active fluid would behave as an isotropic passive one.

III. RESULTS

Let us consider a continuous steady solution $p(z)$ for the polarization field. Then, the steady state solution to Eq. (11) is

$$\rho = \frac{b_2}{b_3} \ln(d + b_3 p^2) + \rho_0, \quad (14)$$

where $d + b_3 p^2 > 0$ is assumed³⁷ and ρ_0 is a constant determined by the condition $L^{-1} \int_0^L \rho dz = 1$. At steady state, the shear stress is uniform across the gap ($\partial_z \sigma = 0$). One can then integrate Eq. (13) for a constant (unknown) σ and obtain the velocity field

$$u = \frac{\sigma}{2\eta}(2z - L) - \xi p + \sqrt{\frac{d}{b_3}} \xi \arctan\left(\sqrt{\frac{b_3}{d}} p\right) - \frac{\beta_\sigma}{\eta} p \rho, \quad (15)$$

with

$$\xi = \frac{b_2}{\eta b_3} (mb_1 + md - 2\beta_\sigma). \quad (16)$$

The (macroscopic) steady-state flow curve $\sigma = f(\dot{\gamma})$, as would be measured by using a rheometer, is then obtained from Eq. (15) by satisfying the no-slip boundary conditions at the moving walls. One finds

$$\sigma = \eta \dot{\gamma} + \sigma_0, \quad (17)$$

with

$$\sigma_0 = \frac{2\eta}{L} \left\{ \xi p_{eq} - \sqrt{\frac{d}{b_3}} \xi \arctan\left(\sqrt{\frac{b_3}{d}} p_{eq}\right) + \frac{\beta_\sigma}{\eta} p_{eq} \left[\frac{b_2}{b_3} \ln(d + b_3 p_{eq}^2) + \rho_0 \right] \right\}. \quad (18)$$

The slope $d\sigma/d\dot{\gamma}$ is simply the fluid viscosity η , as for an isotropic passive fluid; however, the stress at a zero strain rate, σ_0 , is not zero: The active LC has a yield stress and effectively behaves, from a




FIG. 2. The effect of system size: (a) polarization magnitude field, (b) density field, (c) velocity field, and (d) deviation from asymptotic results in the stress at a zero strain rate (inset: stress). Analytical profiles and stress were obtained using Eq. (19) for $L \gg 1$ and Eq. (20) for $L \ll 1$ (here, we set for simplicity $-a_2/K = 1$). The parameters are $a_2 = -1$, $a_4 = 1$, $K = 1$, $\eta = 1$, $b_{1,2,3} = 0.1$, $m = -1$, $d = 1$, $\beta_\sigma = -1$, and $\dot{\gamma} = 0$ (arbitrary units).

rheological point of view, in a similar way to a Bingham fluid.³⁸ Indeed, gradients in p induce reversible and active contributions to the stress which exist independently of the applied strain rate. The converse is also true: A non-zero macroscopic strain rate is required in order to maintain a zero stress. Moreover, σ_0 can be of either sign, meaning that the apparent viscosity, defined as $\sigma/\dot{\gamma}$, can be negative (while passive contributions can only add to σ , active contributions $\sim\beta_\sigma$ can either add or subtract to σ).

There exist, as far as we know, two limiting cases where an explicit steady state analytical solution to Eq. (10) can be written. In the limit $-a_2L^2/K \gg 1$, a good approximation for p is the solution for an infinite system³⁹

$$p = p_{\text{eq}} \tanh \left[\sqrt{\frac{-a_2}{2K}} (z - z_i) \right], \quad (19)$$

where z_i is the (undetermined) location of the interface (whose thickness decreases with increasing $-a_2/K$) between two polar phases pointing in opposite directions. This profile results in depletion (or accumulation, depending on the sign of b_2) of ρ localized at the interface and in a non-uniform, non-monotonic velocity profile.

In the opposite limit $-a_2L^2/K \ll 1$, the solution for the polarization magnitude can be approximated by a linear profile

$$p = -p_{\text{eq}} + \frac{2p_{\text{eq}}}{L}z. \quad (20)$$

Note however that this latter limit may require that the width L be comparable to the active particle size, for which the validity of the hydrodynamic equations is in question and is mostly of interest here as a bound.

In addition to these limiting cases, Eqs. (10), (11), and (13) were also solved numerically. Our algorithm is based on second-order implicit finite difference schemes (Crank-Nicolson scheme for time integration and centered schemes for spatial discretization) with adaptive time-stepping. Time-dependent equations were solved to steady-state, starting from a linear profile for p and a uniform ρ . A comparison between the asymptotic solutions and the numerical ones is shown in Fig. 2, together with additional results for an intermediate value of $-a_2L^2/K$. The estimate of σ_0 obtained from asymptotic expressions is remarkably accurate (deviation not greater than 1%) even for $-a_2L^2/K = O(1)$.

IV. DISCUSSION AND CONCLUSIONS

We considered the minimal problem of a one-dimensional sheared active LC under confinement, with a uniform orientation of the polarization field, focusing on the effect of varying its signed magnitude p . Our analysis thereby complements prior studies of the same system which allowed gradients in the orientation field, while keeping the magnitude of liquid crystalline order constant.^{29,30,40}

As the dynamics of p is not coupled to that of the density or the velocity, the uniform equilibrium solution is always stable and gradients in p must be generated through boundary conditions. Here, we imposed p to be of equal magnitude and of opposite signs at the walls. Such asymmetric polarization at the boundaries could be realized experimentally through manipulation of the surface chemistry or architecture.⁴¹⁻⁴⁵

The case of variable orientation leads to a rich phenomenology, including a spontaneous transition to a flowing state in the absence of external driving,⁴⁰ and the existence of non-monotonic stress vs. strain rate flow curves.^{29,30} By contrast, the case of variable polarization studied here does not yield such unusual mechanical properties: The relationship between the stress and the macroscopic strain rate is linear and, for a nematic active LC, would not differ from that for an isotropic fluid. For a polar active LC though, there exist elastic and active contributions to the total stress in addition to the viscous one, and the flow curve does not pass through the origin. This indicates that macroscopic stresses are present in the uniformly aligned polar active LC even in the absence of external driving.

One of the advantages of the simple configuration considered here lies in the fact that analytical solutions can be explicitly obtained. We hope that these solutions will provide insight into the role played by the gradients of liquid crystalline order and could be used as a starting point and benchmark reference for numerical work on sheared active polar LCs, where both the magnitude and direction of the LC order parameter vary.³²⁻³⁴

ACKNOWLEDGMENTS

Part of this work was funded by a Leverhulme Trust Research Project (Grant No. RPG-2016-147). A.P.T. acknowledges a University of Bristol undergraduate summer bursary. T.B.L. acknowledges support of BrisSynBio, a BBSRC/EPSRC Advanced Synthetic Biology Research Centre (Grant No. BB/L01386X/1).

REFERENCES

- C. Dombrowski, L. Cisneros, S. Chatkaew, R. E. Goldstein, and J. O. Kessler, *Phys. Rev. Lett.* **93**, 098103 (2004).
- H. P. Zhang, A. Be'er, E.-L. Florin, and H. L. Swinney, *Proc. Natl. Acad. Sci. U. S. A.* **107**, 13626 (2010).
- J. Gachelin, A. Rousselet, A. Lindner, and E. Clement, *New J. Phys.* **16**, 025003 (2014).
- C. Chen, S. Liu, X.-Q. Shi, H. Chaté, and Y. Wu, *Nature* **542**, 210 (2017).
- V. Schaller, C. Weber, C. Semmrich, E. Frey, and A. R. Bausch, *Nature* **467**, 73 (2010).
- T. Sanchez, D. T. N. Chen, S. J. DeCamp, M. Heymann, and Z. Dogic, *Nature* **491**, 431 (2012).
- K.-T. Wu, J. B. Hishamunda, D. T. N. Chen, S. J. DeCamp, Y.-W. Chang, A. Fernández-Nieves, S. Fraden, and Z. Dogic, *Science* **355**, eaal1979 (2017).
- A. Bricard, J.-B. Caussin, N. Desreumaux, O. Dauchot, and D. Bartolo, *Nature* **503**, 95 (2013).
- I. Theurkauff, C. Cottin-Bizonne, J. Palacci, C. Ybert, and L. Bocquet, *Phys. Rev. Lett.* **108**, 268303 (2012).
- S. Ramaswamy, *Annu. Rev. Condens. Matter Phys.* **1**, 323 (2010).
- M. C. Marchetti, J. F. Joanny, S. Ramaswamy, T. B. Liverpool, J. Prost, M. Rao, and R. A. Simha, *Rev. Mod. Phys.* **85**, 1143 (2013).
- J. Prost, F. Jülicher, and J.-F. Joanny, *Nat. Phys.* **11**, 111 (2015).
- D. Saintillan and M. J. Shelley, *C. R. Phys.* **14**, 497 (2013).
- F. Jülicher, S. W. Grill, and G. Salbreux, *Rep. Prog. Phys.* **81**, 076601 (2018).
- P. G. de Gennes and J. Prost, *The Physics of Liquid Crystals*, 2nd ed. (Oxford University Press, Oxford, 1993).
- A. Sokolov and I. S. Aranson, *Phys. Rev. Lett.* **103**, 148101 (2009).
- S. Rafai, L. Jibuti, and P. Peyla, *Phys. Rev. Lett.* **104**, 098102 (2010).
- J. Gachelin, G. Miño, H. Berthet, A. Lindner, A. Rousselet, and É. Clément, *Phys. Rev. Lett.* **110**, 268103 (2013).

- ¹⁹M. Mussler, S. Rafai, P. Peyla, and C. Wagner, *Europhys. Lett.* **101**, 54004 (2013).
- ²⁰H. M. López, J. Gachelin, C. Douarche, H. Auradou, and E. Clément, *Phys. Rev. Lett.* **115**, 028301 (2015).
- ²¹D. Saintillan, *Annu. Rev. Fluid Mech.* **50**, 563 (2018).
- ²²S. Guo, D. Samanta, Y. Peng, X. Xu, and X. Cheng, *Proc. Natl. Acad. Sci. U. S. A.* **115**, 7212 (2018).
- ²³Y. Hatwalne, S. Ramaswamy, M. Rao, and R. A. Simha, *Phys. Rev. Lett.* **92**, 118101 (2004).
- ²⁴T. B. Liverpool and M. C. Marchetti, *Phys. Rev. Lett.* **97**, 268101 (2006).
- ²⁵B. M. Haines, A. Sokolov, I. S. Aranson, L. Berlyand, and D. A. Karpeev, *Phys. Rev. E* **80**, 041922 (2009).
- ²⁶D. Saintillan, *Exp. Mech.* **50**, 1275 (2010).
- ²⁷D. Saintillan, *Phys. Rev. E* **81**, 056307 (2010).
- ²⁸S. D. Ryan, B. M. Haines, L. Berlyand, F. Ziebert, and I. S. Aranson, *Phys. Rev. E* **83**, 050904 (2011).
- ²⁹A. Loisy, J. Eggers, and T. B. Liverpool, *Phys. Rev. Lett.* **121**, 018001 (2018).
- ³⁰L. Giomi, T. B. Liverpool, and M. C. Marchetti, *Phys. Rev. E* **81**, 051908 (2010).
- ³¹S. Fürthauer, M. Neef, S. W. Grill, K. Kruse, and F. Jülicher, *New J. Phys.* **14**, 023001 (2012).
- ³²D. Marenduzzo, E. Orlandini, and J. M. Yeomans, *Phys. Rev. Lett.* **98**, 118102 (2007).
- ³³M. E. Cates, S. M. Fielding, D. Marenduzzo, E. Orlandini, and J. M. Yeomans, *Phys. Rev. Lett.* **101**, 068102 (2008).
- ³⁴S. M. Fielding, D. Marenduzzo, and M. E. Cates, *Phys. Rev. E* **83**, 041910 (2011).
- ³⁵E. Tjhung, M. E. Cates, and D. Marenduzzo, *Soft Matter* **7**, 7453 (2011).
- ³⁶W. Kung, M. C. Marchetti, and K. Saunders, *Phys. Rev. E* **73**, 031708 (2006).
- ³⁷Otherwise this would be equivalent to a negative diffusion coefficient in the density equation, meaning that fluctuations around the equilibrium density would not remain small and that stabilizing higher order terms would need to be included in our model.
- ³⁸E. C. Bingham, *Fluidity and Plasticity* (McGraw-Hill, 1922).
- ³⁹G. F. Mazenko, *Fluctuations, Order, and Defects* (John Wiley & Sons, Hoboken, NJ, 2003).
- ⁴⁰R. Voituriez, J. F. Joanny, and J. Prost, *Europhys. Lett.* **70**, 404 (2005).
- ⁴¹N. Koumakis, C. Maggi, and R. Di Leonardo, *Soft Matter* **10**, 5695 (2014).
- ⁴²O. Sipos, K. Nagy, R. Di Leonardo, and P. Galajda, *Phys. Rev. Lett.* **114**, 258104 (2015).
- ⁴³Y. Yuan and Y. Zhang, *Nanomed.: Nanotechnol., Biol. Med.* **13**, 2199 (2017).
- ⁴⁴A. Muñoz-Bonilla, R. Cuervo-Rodríguez, F. López-Fabal, J. L. Gómez-Garcés, and M. Fernández-García, *Materials* **11**, 1266 (2018).
- ⁴⁵J. Hasan, R. J. Crawford, and E. P. Ivanova, *Trends Biotechnol.* **31**, 295 (2013).