

# QUANTUM COMPUTATION

## Exercise sheet 6

Ashley Montanaro, University of Bristol

ashley.montanaro@bristol.ac.uk

1. **Shor's 9 qubit code.** Imagine we encode the state  $|\psi\rangle = \alpha|0\rangle + \beta|1\rangle$  using Shor's 9 qubit code, and then an  $X$  error occurs on the 8th qubit of the encoded state  $|E(\psi)\rangle$ .

(a) Write down the state following the error.

**Answer:**

$$\frac{1}{2\sqrt{2}}(\alpha(|000\rangle+|111\rangle)(|000\rangle+|111\rangle)(|010\rangle+|101\rangle)+\beta(|000\rangle-|111\rangle)(|000\rangle-|111\rangle)(|010\rangle-|101\rangle)).$$

- (b) We now decode the encoded state, starting by applying the bit-flip code decoding algorithm. What are the syndromes returned by the measurements in the algorithm?

**Answer:** Using the table in the lecture notes, the syndromes are 00, 00, 10.

- (c) Now imagine that  $|E(\psi)\rangle$  is affected by two  $X$  errors, on the 7th and 8th qubits. What are the syndromes returned this time? What state does the decoding algorithm output?

**Answer:** Now the syndromes are 00, 00, 01. The decoding algorithm thus thinks there has been an  $X$  error on the 9th qubit. So it "corrects" this by applying an  $X$  operation on this qubit, to give the state

$$\frac{1}{2\sqrt{2}}(\alpha(|000\rangle+|111\rangle)(|000\rangle+|111\rangle)(|000\rangle+|111\rangle)-\beta(|000\rangle-|111\rangle)(|000\rangle-|111\rangle)(|000\rangle-|111\rangle)).$$

Note that  $\beta$  now has a minus sign in front of it. After the bit-flip decoding, we are left with  $\alpha|+++ \rangle - \beta|--- \rangle$ , which is then decoded to  $\alpha|0\rangle - \beta|1\rangle$ .

- (d) Which patterns of  $X$  errors are corrected by Shor's 9 qubit code?

**Answer:** If there is at most one  $X$  error in each block of 3 qubits, these will be corrected properly. We have just seen that, if two errors occur in one block, the sign of  $\beta$  will be flipped, but the state is not otherwise affected; a similar argument holds for 3 errors in one block. So the output state will be correct if the number of blocks in which at least two errors occur is even (as then  $\beta$  will eventually be left unchanged).