

ICTP Summer School on Dynamical Systems

Renormalization in entropy zero dynamics (week 2)

Homework 5

Exercise C1 Let $d = 2$ and let T be a 2-IET with initial lengths $(\alpha, 1 - \alpha)$ where $\alpha \in (0, 1)$ irrational.

- (a) Apply Rauzy-Veech induction to T and verify that if we set

$$k := \begin{cases} \left[\frac{1-\alpha}{\alpha} \right] & \text{if } \alpha < 1/2 \\ \left[\frac{\alpha}{1-\alpha} \right] & \text{if } \alpha > 1/2 \end{cases}.$$

(where $[x]$ denotes the integer part of x), Rauzy-Veech induction starts with either k type top (if $\alpha < 1/2$) or k type top (if $\alpha > 1/2$) moves.

- (b) For any $n \in \mathbb{N}$, if $\lambda_A^{(n)}$ and $\lambda_B^{(n)}$ are the lengths of the intervals exchanged by $T^{(n)}$ (n^{th} induced map in the induction), set $\alpha_n \in [0, 1]$ to be given by

$$\alpha_n := \begin{cases} \frac{\lambda_A^{(n)}}{\lambda_B^{(n)}} & \text{if } \lambda_A^{(n)} < \lambda_B^{(n)} \\ \frac{\lambda_B^{(n)}}{\lambda_A^{(n)}} & \text{if } \lambda_B^{(n)} < \lambda_A^{(n)} \end{cases}$$

- . Show that $\alpha_n = \mathcal{F}^n(\alpha)$ where $\mathcal{F}(\alpha)$ is the Farey map given by

$$F(x) = \begin{cases} \frac{1-x}{x} & \text{if } x > \frac{1}{2} \\ \frac{x}{1-x} & \text{if } x \leq \frac{1}{2}. \end{cases}$$

[Recalling Exercise 3.3 of Week 1, this also shows that the acceleration of Rauzy-Veech induction obtained by doing in one go all bottom or all top steps, plus the next one of a different type, corresponds to the Gauss map in the special case $d = 2$. This acceleration is known as *Zorich induction*. One can show that Zorich induction preserves a *finite* invariant measure on the space of d -IETs, while Rauzy-Veech induction preserves a *infinite* invariant measure (as the Gauss map preserves the Gauss measure, which is finite, while the Farey map \mathcal{F} preserves a measure with density $1/x$ which is infinite.)]

TURN PAGE for next Exercise

Exercise C2 (Renormalization for Birkhoff sums) Let T be a d -IET (with π irreducible and $(\lambda_\alpha)_{\alpha \in \mathcal{A}}$ rationally independent) and let $f : I \rightarrow \mathbb{R}$. Let $I^{(n)}$ be the inducing intervals for Rauzy Veech induction and define the sequence of induced functions by

$$f^{(n)}(x) := \sum_{i=0}^{h_\alpha^{(n)}-1} f(T^i(x)) \quad \text{if } x \in I_\alpha^{(n)}.$$


[This function associate to a point in the base $I^{(n)}$ the Birkhoff sum of the function f along the tower, i.e. along the piece of the orbit that goes from the base point up to the top of the tower to which it belongs. They are also called *special Birkhoff sums*.]

- (a) Show that if f is constant on $I_\alpha^{(0)}$ for each $\alpha \in \mathcal{A}$, then for every $n \in \mathbb{N}$ also $f_\alpha^{(n)}$ is constant on $I_\alpha^{(n)}$ for each $\alpha \in \mathcal{A}$.
- (b) Let $f_\alpha^{(n)}$ be the value of $f^{(n)}$ on any point $x \in I_\alpha^{(n)}$ (so that by (a) $f^{(n)} \equiv f_\alpha^{(n)}$ on $I_\alpha^{(n)}$) and let $f^{(n)}$ be the vector with coordinates $f_\alpha^{(n)}$, $\alpha \in \mathcal{A}$. Show that

$$f^{(n+1)} = A_n^t f^{(n)},$$

where A_n is the matrix that transforms lengths vectors in Rauzy-Veech induction (i.e. $f^{(n)}$ transforms as the tower heights/return times vector $h^{(n)}$ (compare with Homework 4 of this week).

[*Hint*: you might want to use cutting and stacking of towers, see e.g. the figure.]



- (c) Deduce that the entry $(A_n^t \cdots A_0^t)_{\alpha\beta}$ of the matrix product $A_n^t \cdots A_0^t$ gives the number of time the orbit under T of the interval $I_\beta^{(n)}$ visits the interval $I_\alpha^{(0)}$ until the first return time $h_\beta^{(n)}$ to $I^{(n)}$, or equivalently the number of floors of the tower $H_\beta^{(n)}$ which are contained in $I_\alpha^{(0)}$.
- (d) Assume that T is periodic under renormalization, i.e. $T^{(n)}$ is equal to T up to rescaling the interval $I^{(n)}$ to unit length (T in this case is self-similar). Show that we can decompose

$$\mathbb{R}^d = E^- \oplus E^+$$

such that for every $(f_\alpha)_{\alpha \in \mathcal{A}} \in E^-$, special Birkhoff sums $f^{(n)}$ are uniformly bounded for each n , while for every $(f_\alpha)_{\alpha \in \mathcal{A}} \notin E^-$, $\|f^{(n)}\|_\infty$ grows exponentially.

[The first part can be used to show that, among piecewise constant functions, coboundaries, i.e. functions that can be written as $f = g \circ T - g$ for some $g : I \rightarrow \mathbb{R}$, can only be found among functions f given by $(f_\alpha)_{\alpha \in \mathcal{A}} \in E^-$. Using the second part, one can show that Birkhoff sums of mean zero functions f display polynomial deviations of ergodic averages, i.e. for a.e. $x \in [0, 1)$ $S_n f(x) = o(n)$ (by ergodicity) but there exists $0 < \gamma < 1$ such that $S_n f(x) \geq cn^\gamma$ infinitely often.]