

MATH20901 Multivariable Calculus: Problems 5 ¹

1. Verify Stokes' theorem for the (hemispheric) surface $|\mathbf{r}| = 3, z \geq 0$ and the vector field $\mathbf{v}(\mathbf{r}) = (y, -x, 0)$.
2. If there exists a scalar function $f(x, y)$ such that a vector field $\mathbf{F} = \nabla f$ show, using Green's theorem in the plane, that $\int_C \mathbf{F} \cdot d\mathbf{r} = 0$ where C is a closed curve.
3. Deduce the value of integral in Problem Sheet 4 Q4 when the square is rotated in the (x, y) -plane by an arbitrary angle about the vertex at $(0, 0)$.
4. (a) Compute

$$\int_S \mathbf{F} \cdot d\mathbf{S}$$

when $\mathbf{F} = (x, y, z^4)$ and S is the conical surface formed by $z = \sqrt{x^2 + y^2}$ for $0 < z < 1$; the unit normal to the surface is directed away from the axis of the cone.

- (b) Find

$$\int_V \nabla \cdot \mathbf{F} dV$$

where V is the volume inside the cone defined above.

- (c) Without doing any further computations, deduce

$$\int_{x^2+y^2 < 1, z=1} \mathbf{F} \cdot d\mathbf{S}.$$

Check your answer.

5. Evaluate $\int_{\partial V} \mathbf{F} \cdot d\mathbf{S}$, where $\mathbf{F}(x, y, z) = x\hat{\mathbf{x}} + y\hat{\mathbf{y}} - z\hat{\mathbf{z}}$ and V is the unit cube in the first octant. Perform the calculation directly and check by using the divergence theorem.
6. On Problem Sheet 2, Q4(b) it was shown that $\Delta\left(\frac{1}{r}\right) = 0$ for $r \neq 0$. Let V be a sphere of arbitrary non-zero radius centred on the origin and S its enclosing surface. Use the divergence theorem to show that

$$\int_V \Delta\left(\frac{1}{r}\right) dV = -4\pi.$$

How do you explain this ?

7. Gauss' Law relates the flux of an electric field $\mathbf{E}(\mathbf{r})$ through a surface S to the density of charges in V , within S , $\rho(\mathbf{r})$ via

$$\int_S \mathbf{E} \cdot d\mathbf{S} = 4\pi \int_V \rho dV = Q$$

where Q is the net charge.

¹©University of Bristol 2018

This material is copyright of the University of Bristol unless explicitly stated. It is provided exclusively for educational purposes at the University of Bristol and is to be downloaded or copied for your private study only.

(a) Show that (one of Maxwell's equations)

$$\nabla \cdot \mathbf{E} = 4\pi\rho, \quad \text{at all points in space}$$

(b) Compute Q when $\mathbf{E} = (x, y, z)$ and V is the cuboid with vertices $(\pm 1, \pm 1, \pm 1)$. (Hint: use the easiest method you can).

8. We are all happy with the idea that volume of a region $V \subset \mathbb{R}^3$ can be defined as $\int_V dx dy dz$. Show that an alternative way of calculating it is

$$\frac{1}{3} \int_S \mathbf{r} \cdot \hat{\mathbf{n}} dS$$

where S is the surface enclosing V and $\hat{\mathbf{n}}$ the unit outward normal to S .

Confirm your answer by applying the new formula directly to a sphere of radius a .

9. In a style similar to Q8, can you think of a way of computing surface areas of 3D objects using integrals over the volume ?
10. Use Green's identity Identity to show, for two scalar fields $u(\mathbf{r})$, $v(\mathbf{r})$ that, if $\Delta u = 0$ in a volume V bounded by a surface S , upon which $v = 0$,

$$\int_V \nabla u \cdot \nabla v dV = 0$$