UNIVERSITY OF BRISTOL

Examination for the Degrees of B.Sc. and M.Sci. (Levels 3 and M)

STATISTICAL MECHANICS 3 MATH 34300 (Paper Codes MATH-34300)

May-June 2011, 2 hours 30 minutes

This paper contains **five** questions A candidate's **FOUR** best answers will be used for assessment. Calculators are **not** permitted in this examination.

You may wish to use Stirling's formula which for $N \gg 1$, gives

 $\ln N! \approx N \ln N - N \quad .$

For small x,

$$\cosh x = 1 + \frac{x^2}{2} + \frac{x^4}{4!} + \dots ,$$

$$\sinh x = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots ,$$

$$\coth x = \frac{1}{x} + \frac{x}{3} - \frac{x^3}{45} + \dots$$

Do not turn over until instructed.

Cont...

,

- 1. (a) (4 marks) Consider a gas of N particles in a volume V.
 - i. At pressure p, what is the differential work done in expanding an infinitesimal volume dV?
 - ii. Use the 2nd law of thermodynamics to obtain expressions for

$$\left(\frac{\partial S}{\partial V}\right)_{E,N}$$
 and $\left(\frac{\partial E}{\partial V}\right)_{S,N}$,

where S is the entropy and E the internal energy of the gas.

(b) (8 marks)

Use the fact that the entropy, S(E, V, N), of a gas composed of N molecules with energy E, volume V is *extensive* to show that

$$S = E\left(\frac{\partial S}{\partial E}\right)_{V,N} + V\left(\frac{\partial S}{\partial V}\right)_{E,N} + N\left(\frac{\partial S}{\partial N}\right)_{E,V}$$

and hence that $E = TS - pV + \mu N$

- (c) Consider using a diatomic ideal gas composed of N molecules as the working substance of an engine whose reversible cycle $(1\rightarrow 2\rightarrow 3\rightarrow 1)$ consists of three quasistatic steps: $1\rightarrow 2$ is isobaric expansion (at constant pressure), $2\rightarrow 3$ is a constant volume process (isochoric), $3\rightarrow 1$ is isothermal compression (at constant temperature). The temperatures T_1 and T_2 at points 1 and 2 in the cycle are given.
 - i. (3marks) Draw this cycle in p V coordinates.
 - ii. (6 marks) For each step compute the work done by the gas and the heat added to the gas. Identify which steps have heat flowing into the gas.
 - iii. (4 marks) What is the efficiency of the cycle in terms of T_1 and T_2 ?

- 2. (a) Consider a system in a macrostate of fixed E in the microcanonical ensemble.
 - i. (2 marks) What is the probability of the system being in a particular microstate?
 - ii. (3 marks) Write down an expression for the entropy S(E).
 - iii. (5 marks) Use the microcanonical ensemble to show that if two macroscopic systems with fixed energies E_1 and E_2 , respectively are brought into contact, they will be in equilibrium only if

$$\frac{\partial S_1}{\partial E_1} = \frac{\partial S_2}{\partial E_2}$$

(b) A simplified model of a magnet in a magnetic field **B** is a collection of *N* distinguishable spins which can each be characterised by a magnetic moment, a *three* dimensional vector of fixed magnitude μ and arbitrary orientation, \mathbf{S}_i , $i \in \{1, \ldots, N\}$. Due to its magnetic moment each 'spin' has lower energy when aligned with the field. The system is thus described by a Hamiltonian

$$\mathcal{H} = -\mu \sum_{i=1}^{N} \mathbf{B} \cdot \mathbf{S}_{i} , \quad |\mathbf{S}_{i}| = 1 ,$$

and the magnetization is given by $\mathbf{M} = \mu \sum_{i=1}^{N} \mathbf{S}_i$. Consider such a 'magnet' at a fixed temperature, T with the magnetic field $\mathbf{B} = B \hat{\mathbf{z}}$ (oriented along the z-axis).

- i. (4 marks) Which ensemble is appropriate for this system ? How can a microstate be characterised?
- ii. (3 marks) What is the probability of being in a particular microstate ?
- iii. (8 marks) Calculate the Helmholtz free energy and the average magnetization, $\langle \mathbf{M} \rangle$. Hence obtain an expression for the susceptibility, $\chi(T)$ defined as

$$\chi(T) = \left(\frac{\partial}{\partial \mathbf{B}} \cdot \langle \mathbf{M} \rangle\right)_{B=0} \; .$$

- 3. (a) Consider a dilute gas of N particles in three dimensions with positions $\mathbf{q}_i(t)$ and momenta $\mathbf{p}_i(t)$, $(1 \le i \le N)$, which satisfy Hamilton's equations of motion. Its microstate is defined by the 6N-dimensional phase space (\mathbf{p}, \mathbf{q}) where $\mathbf{p} = (\mathbf{p}_1, \cdots, \mathbf{p}_N), \mathbf{q} = (\mathbf{q}_1, \cdots, \mathbf{q}_N)$ with Hamiltonian $\mathcal{H}(\mathbf{p}, \mathbf{q})$.
 - i. (5 marks) What is the phase space density $\rho(\mathbf{p}, \mathbf{q}, t)$?
 - ii. (10 marks) Show that a consequence of Liouville's theorem is that the phase space density, $\rho(\mathbf{p}, \mathbf{q}, t)$ satisfies the equation

$$\frac{d\rho}{dt} = \frac{\partial\rho}{\partial t} + \{\rho, \mathcal{H}\} = 0 \tag{1}$$

and explain the meaning of the **Poisson bracket** $\{A, B\}$.

(b) (10 marks) Non Hamiltonian dynamics in molecular dynamics simulations: consider a system of N particles in three dimensions with Hamiltonian, \mathcal{H} :

$$\mathcal{H}(\{\mathbf{p}_i, \mathbf{q}_i\}) = \sum_{i=1}^N \frac{1}{2m} |\mathbf{p}_i|^2 + \sum_{i \neq j} U(\mathbf{q}_i - \mathbf{q}_j) ,$$

where the equation of motion is modified by a friction term to

$$\dot{\mathbf{q}}_i = rac{\partial \mathcal{H}}{\partial \mathbf{p}_i} \;\;;\;\;\; \dot{\mathbf{p}}_i = -rac{\partial \mathcal{H}}{\partial \mathbf{q}_i} - \lambda \mathbf{p}_i$$

Given that $\lambda(\mathbf{p})$ is function of momenta only, show that the Liouville theorem can be written in the form

$$\frac{\partial \rho}{\partial t} + \{\rho, \mathcal{H}\} = -C\lambda\rho + \boldsymbol{D} \cdot \frac{\partial \rho}{\partial \boldsymbol{p}} ,$$

and give the value of C, D in terms of λ and p.

4. (a) (4 marks)

Write down expressions for the classical and quantum partition functions in the *canonical ensemble* for a set of N particles with momentum p_i and position q_i and Hamiltonian $\mathcal{H}(\{p_i, q_i\})$. Explain why Planck's constant appears in the classical formula.

(b) (8 marks)

The Hamiltonian for a classical harmonic oscillator of frequency ω is

$$\mathcal{H}\left(\{q,p\}\right) = \frac{p^2}{2m} + \frac{m\omega^2 q^2}{2} .$$

Calculate the free energy ${\cal F}$, entropy ${\cal S}$ and internal energy ${\cal E}.$

(c) (8 marks)

The energy levels of a quantum harmonic oscillator of frequency ω are given by

$$\mathcal{E}_n = \hbar \omega \left(n + \frac{1}{2} \right) , \ n = 0, 1, 2, \cdots$$

Calculate the free energy F, entropy S and internal energy E.

(d) (5 marks)

Compare and contrast the specific heat capacity obtained from the classical and quantum calculations (it may be helpful to sketch c(T)). Comment on what happens as $T \to 0$ and at high temperatures.

- 5. (a) Consider a *classical* system in the grand canonical ensemble.
 - i. (4 marks) Write down the expression for the *classical* grand partition function. How is this related to the grand potential?
 - ii. (6 marks) Calculate the mean number of particles $\langle N \rangle$ and the mean square fluctuations $\langle \Delta N^2 \rangle = \langle N^2 \rangle \langle N \rangle^2$ and hence obtain the relative mean-square fluctuation $\frac{\langle \Delta N^2 \rangle}{\langle N \rangle^2}$ in the thermodynamic limit.
 - (b) An ideal gas of classical *indistinguishable* particles of mass m is in a uniform attractive potential $U(\mathbf{r}) = -\varepsilon$ where $\varepsilon > 0$ in *three* dimensions (i.e. a particle at position \mathbf{q}_i has a potential energy $U(\mathbf{q}_i)$). The gas is in contact with a reservoir at fixed chemical potential μ and temperature T and is confined to a rectangular box of height h and base area A.
 - i. (3 marks) Write down the Hamiltonian of the system.
 - ii. (5 marks) Evaluate the grand partition function \mathcal{Z} and hence the grand potential, \mathcal{G} of the system.
 - iii. (7 marks) Calculate the pressure, p and density, ρ and hence obtain a relationship between ρ , p and T.

End of examination.