

# Waring's Problem: A Survey

R. C. Vaughan<sup>1</sup> and T. D. Wooley<sup>2</sup>

## 1 The Classical Waring Problem

*“Omnis integer numerus vel est cubus, vel e duobus, tribus, 4, 5, 6, 7, 8, vel novem cubis compositus, est etiam quadrato-quadratus vel e duobus, tribus, &c. usque ad novemdecim compositus, & sic deinceps.”*

Waring [150, pp. 204-5].

“Every integer is a cube or the sum of two, three, ... nine cubes; every integer is also the square of a square, or the sum of up to nineteen such; and so forth.”

Waring [152, p. 336].

It is presumed that by this, in modern notation, Waring meant that for every  $k \geq 3$  there are numbers  $s$  such that every natural number is the sum of at most  $s$   $k$ -th powers of natural numbers and that the smallest such number  $g(k)$  satisfies  $g(3) = 9$ ,  $g(4) = 19$ .

By the end of the nineteenth century, the existence of  $g(k)$  was known for only a finite number of values of  $k$ . There is an account of this work in Dickson [48], and as far as we have been able to ascertain, by 1909 its existence was known for  $k = 3, 4, 5, 6, 7, 8, 10$ , but not for any larger  $k$  (of course, with the natural extension of the definition of  $g(k)$ , Lagrange proved in 1770 that  $g(2) = 4$ ). However, starting with Hilbert [69], who showed that  $g(k)$  does indeed exist for every  $k$ , the twentieth century has seen an almost complete solution of this problem. Let  $[x]$  denote the greatest integer not exceeding  $x$  and write  $\{x\}$  for  $x - [x]$ . As the result of the work of many mathematicians we now know that

$$g(k) = 2^k + [(3/2)^k] - 2,$$

provided that

$$2^k \{(3/2)^k\} + [(3/2)^k] \leq 2^k. \tag{1.1}$$

If this fails, then

$$g(k) = 2^k + [(3/2)^k] + [(4/3)^k] - \theta$$

---

<sup>1</sup>Research supported by NSF grant DMS-9970632.

<sup>2</sup>Packard Fellow, and supported in part by NSF grant DMS-9970440.

where  $\theta$  is 2 or 3 according as

$$[(4/3)^k][(3/2)^k] + [(4/3)^k] + [(3/2)^k]$$

equals or exceeds  $2^k$ .

The condition (1.1) is known to hold (Kubina & Wunderlich [85]) whenever  $k \leq 471, 600, 000$ , and Mahler [91] has shown that there are at most a finite number of exceptions. To complete the proof for all  $k$  it would suffice to know that  $\{(3/2)^k\} \leq 1 - (3/4)^{k-1}$ . Beukers [3] has shown that whenever  $k > 5,000$  one has  $\{(3/2)^k\} \leq 1 - a^k$ , where  $a = 2^{-0.9} = 0.5358\dots$ , and this has been improved slightly by Dubitskas [49] to  $a = 0.5769\dots$ , so long as  $k$  is sufficiently large (see also Bennett [1] for associated estimates). A problem related to the evaluation of  $g(k)$  now has an almost definitive answer. Let  $g_r(k)$  denote the smallest integer  $s$  with the property that every natural number is the sum of at most  $s$  elements from the set  $\{1^k, r^k, (r+1)^k, \dots\}$ . Then Bennett [2] has shown that for  $4 \leq r \leq (k+1)^{1-1/k} - 1$ , one has  $g_r(k) = r^k + [(1+1/r)^k] - 2$ .

By the way, before turning to the modern form of Waring's problem, it is perhaps worth observing that in the 1782 edition of *Meditationes Algebraicae*, Waring makes an addition:

*“confimilia etiam affirmari possunt (exceptis excipiendis) de eodem numero quantitatum earundem dimensionum.”*

Waring [151, p. 349].

“similar laws may be affirmed (exceptis excipiendis) for the correspondingly defined numbers of quantities of any like degree.”

Waring [152, p. 336].

It would be interesting to know exactly what Waring had in mind. This, taken with some of the observations which immediately follow the remark, suggest that for more general polynomials than the  $k$ -th powers he was aware that some kind of local conditions can play a rôle in determining when representations occur.

## 2 The Modern Problem

The value of  $g(k)$  is determined by the peculiar behaviour of the first three or four  $k$ -th powers. A much more challenging question is the value, for  $k \geq 2$ , of the function  $G(k)$ , the smallest number  $t$  such that every *sufficiently large* number is the sum of at most  $t$   $k$ -th powers of positive integers. The function  $G(k)$  has only been determined for two values of  $k$ , namely  $G(2) = 4$ , by Lagrange in 1770, and  $G(4) = 16$ , by Davenport [30]. The bulk of what is known about  $G(k)$  has been obtained through the medium

of the Hardy-Littlewood method. This has its genesis in a celebrated paper of Hardy and Ramanujan [64] devoted to the partition function. In this paper (section 7.2) there is also a brief discussion about the representation of a natural number as the sum of a fixed number of squares of integers, and there seems little doubt that it is the methods described therein which inspired the later work of Hardy and Littlewood.

Our knowledge concerning the function  $G(k)$  currently leaves much to be desired. If, instead of insisting that all sufficiently large numbers be represented in a prescribed form, one rather asks that *almost all* numbers (in the sense of natural density) be thus represented, then the situation is somewhat improved. Let  $G_1(k)$  denote the smallest number  $u$  such that almost every number  $n$  is the sum of at most  $u$   $k$ -th powers of positive integers. The function  $G_1(k)$  has been determined in five non-trivial instances as follows:

$$\begin{aligned} \text{Davenport [29], } & G_1(3) = 4, \\ \text{Hardy and Littlewood [62], } & G_1(4) = 15, \\ \text{Vaughan [121], } & G_1(8) = 32, \\ \text{Wooley [155], } & G_1(16) = 64, \\ \text{Wooley [155], } & G_1(32) = 128 \end{aligned}$$

(of course, the conclusion  $G_1(2) = 4$  is classical).

### 3 General Upper Bounds for $G(k)$

The first explicit general upper bound for  $G(k)$ , namely

$$G(k) \leq (k-2)2^{k-1} + 5,$$

was obtained by Hardy and Littlewood [61] (in [58] and [59], only the existence of  $G(k)$  is stated, although it is already clear that in principle their method gave an explicit upper bound). In Hardy and Littlewood [62] this is improved to

$$G(k) \leq (k-2)2^{k-2} + k + 5 + [\zeta_k],$$

where

$$\zeta_k = \frac{(k-2)\log 2 - \log k + \log(k-2)}{\log k - \log(k-1)}.$$

There has been considerable activity reducing this upper bound over the years, and Table 1 below presents upper bounds for  $G(k)$  that were probably the best that were known at the time they appeared, at least for

	Vinogradov [136],	$32(k \log k)^2$ ,
	Vinogradov [137][139],	$k^2 \log 4 + (2 - \log 16)k$ ( $k \geq 3$ ),
Vinogradov [135] [138] [140] [143],		$6k \log k + 3k \log 6 + 4k$ ( $k \geq 14$ ),
	Vinogradov [147],	$k(3 \log k + 11)$ ,
	Tong [114],	$k(3 \log k + 9)$ ,
	Jing-Run Chen [24],	$k(3 \log k + 5.2)$ ,
	Vinogradov [148],	$2k(\log k + 2 \log \log k + O(\log \log \log k))$ ,
	Vaughan [124],	$2k(\log k + \log \log k + O(1))$ ,
	Wooley [155],	$k(\log k + \log \log k + O(1))$ .

**Table 1.** General upper bounds for  $G(k)$

$k$  sufficiently large. This list is not exhaustive. In particular, there is a long sequence of papers by Vinogradov between 1934 and 1947, and for further details we refer the reader to the Royal Society obituary of I. M. Vinogradov (see Cassels and Vaughan [23]).

The last entry on this list has been refined further by Wooley [159], and this provides the estimate

$$G(k) \leq k(\log k + \log \log k + 2 + O(\log \log k / \log k))$$

that remains the sharpest available for larger exponents  $k$ .

## 4 Cubes

For small values of  $k$  there are many special variants of the Hardy-Littlewood method that have been developed. However, in the case of cubes, until recently the best upper bounds were obtained by rather different methods that related cubes to quadratic forms, especially sums of squares. Thus Landau [86] had shown that  $G(3) \leq 8$ , and this bound was reduced by Linnik [87][88] to  $G(3) \leq 7$ , with an alternative and simpler proof given by Watson [153]. Only with the advent of refinements to the circle method utilising efficient differencing did it become feasible (Vaughan [119]) to give a proof of the bound  $G(3) \leq 7$  via the Hardy-Littlewood method. Subsequent developments involving the use of smooth numbers (see Vaughan [124][125] and Wooley [158]) have provided a more powerful approach to this problem that, from a practical point of view, is more direct than earlier treatments. Complicated nonetheless, these latter proofs yield much more

information concerning Waring's problem for cubes. We can illustrate the latter observation with two examples which, in the absence of foreseeable progress on the upper bound for  $G(3)$ , provide the problems central to current activity surrounding Waring's problem for cubes.

When  $X$  is a large real number, denote by  $E(X)$  the number of positive integers not exceeding  $X$  that *cannot* be written as a sum of four positive integral cubes. Then the conclusion  $G_1(3) = 4$ , attributed above to Davenport [29], is an immediate consequence of the estimate  $E(X) \ll X^{29/30+\epsilon}$  established in the latter paper. Following subsequent work of Vaughan [119], Brüdern [10][12], and Wooley [158], the sharpest conclusion currently available (see Wooley [162]) shows that  $E(X) \ll X^{1-\beta}$  for any positive number  $\beta$  smaller than

$$(422 - 6\sqrt{2833})/861 = 0.119215\dots$$

It is conjectured that  $G(3) = 4$  (see §10 below), and this would imply that  $E(X) \ll 1$ .

Consider next the density of integers represented as a sum of three positive integral cubes. When  $X$  is a large real number, let  $N(X)$  denote the number of positive integers of the latter type not exceeding  $X$ . It is conjectured that  $N(X) \gg X$ , and following work of Davenport [29][33], Vaughan [118][119], Ringrose [100], Vaughan [124] and Wooley [158], the sharpest currently available conclusion due to Wooley [162] establishes that  $N(X) \gg X^{1-\alpha}$  for any real number  $\alpha$  exceeding

$$(\sqrt{2833} - 43)/123 = 0.083137\dots$$

We remark that, subject to the truth of an unproved Riemann Hypothesis concerning certain Hasse-Weil  $L$ -functions, one has the conditional estimate  $N(X) \gg X^{1-\epsilon}$  due to Hooley [73][74], and Heath-Brown [67]. Unfortunately, the underlying  $L$ -functions are not yet known even to have an analytic continuation inside the critical strip.

We finish our discussion of Waring's problem for cubes by noting that, while Dickson [47] was able to show that 23 and 239 are the only positive integers not represented as the sum of eight cubes of natural numbers, no such conclusion is yet available for sums of seven or fewer cubes (but see McCurley [92] for sums of seven cubes, and Deshouillers, Hennecart and Landreau [44] for sums of four cubes).

## 5 Biquadrates

Davenport's definitive statement that  $G(4) = 16$  is not the end of the story for sums of fourth powers (otherwise known as biquadrates). Let

$G^\#(4)$  denote the least integer  $s_0$  such that whenever  $s \geq s_0$ , and  $n \equiv r \pmod{16}$  for some integer  $r$  with  $1 \leq r \leq s$ , then  $n$  is the sum of at most  $s$  biquadrates. Then Davenport [30] showed that  $G^\#(4) \leq 14$ , and this has been successively reduced by Vaughan [121][124] to  $G^\#(4) \leq 12$ . In an ironic twist of fate, the polynomial identity

$$x^4 + y^4 + (x + y)^4 = 2(x^2 + xy + y^2)^2,$$

reminiscent of identities employed in the nineteenth century, has recently been utilised to make yet further progress. Thus, when  $n \equiv r \pmod{16}$  for some integer  $r$  with  $1 \leq r \leq 10$ , Kawada and Wooley [82] have shown that  $n$  is the sum of at most 11 biquadrates. This and allied identities have also permitted the proof of an effective version of Davenport's celebrated theorem. Thus, as a consequence of work of Deshouillers, Hennecart and Landreau [45] and Deshouillers, Kawada and Wooley [46], it is now known that all integers exceeding 13,792 may be written as the sum of at most sixteen biquadrates. A detailed history of Waring's problem for biquadrates is provided in Deshouillers, Hennecart, Kawada, Landreau and Wooley [43].

## 6 Upper Bounds for $G(k)$ when $5 \leq k \leq 20$

Although we have insufficient space to permit a comprehensive account of the historical evolution of available upper bounds for  $G(k)$  for smaller values of  $k$ , in Table 2 we have recorded many of the key developments, concentrating on the past twenty-five years. Each row in this table presents the best upper bound known for  $G(k)$ , for the indicated values of  $k$ , at the time of publication of the cited work. We note that the claimed bound  $G(7) \leq 52$  of Sambasiva Rao [99] is based on an arithmetical error, and hence we have attributed the bound  $G(7) \leq 53$  parenthetically to Davenport's methods [28] [32]. Also, it is worth remarking that the work of Vaughan and Wooley [130] [131] and [132] appeared in print in an order reversed from its chronological development (indeed, this work was first announced in 1991). The bounds parenthetically attributed to Vaughan and Wooley [132] follow directly from the methods therein, and were announced on that occasion, though details (with additional refinements) appeared only in Vaughan and Wooley [134]. Meanwhile, the bounds recorded in Wooley [159] were an immediate consequence of the methods of Wooley [155] combined with the new estimates for smooth Weyl sums obtained in the former work (no attempt was made therein to exploit the methods of Vaughan and Wooley [132]).

	k	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
Hardy & Littlewood [62]		41	87	192	425	949	2113										
James [81]		35		164		824											
Heilbronn [68]					164	190	217	244	272	300	329	359	388	418	449	480	511
Estermann [52]		29	42	59	78	101	125	153	184	217	253	292	333	377	424	474	
Hua [76]		28															
Davenport [28][32]		23	36	(53)													
Narasimhamurti [96]					73	99	122										
Chen [24]							121										
Cook [26]						96											
Vaughan [115]						91	107	122	137	153	168	184	200	216			
Thanigasalam [107]						90	106	121	136	152	167	183	199	215	231	248	264
Thanigasalam [108]						88	104	119	134	150	165	181	197	213	229	245	262
Thanigasalam [109]				50	68	87	103										
Vaughan [120][121]		21	31	45	62	82											
Vaughan [124][126]		19	29	41	57	75	93	109	125	141	156	171	187	202	217	232	248
Bridern [11]		18															
Vaughan & Wooley [129]		18	28				92	108	124	139	153	168	183	198	213	228	243
Wooley [155]			27	36	47	55	63	70	79	87	95	103	112	120	129	138	146
Wooley [159]						62		78	86	94	102	110	118	127	135	144	
Vaughan & Wooley [132]		17	25	33	43	51	(59)	(67)	(76)	(84)	(92)	(100)					
Vaughan & Wooley [130]					42												
Vaughan & Wooley [131]			24														
Meng [94]																	143
Vaughan & Wooley [134]						50	59	67	76	84	92	100	109	117	125	134	142
Conjectured		6	9	8	32	13	12	12	16	14	15	16	64	18	27	20	25

Table 2. Upper bounds for  $G(k)$  when  $5 \leq k \leq 20$

## 7 The Hardy-Littlewood Method

Practically all of the above conclusions have been obtained via the Hardy-Littlewood method. Here is a quick introduction. Let  $n$  be a large natural number, and write  $P = n^{1/k}$  and

$$f(\alpha) = \sum_{x \leq P} e(\alpha x^k)$$

(here we follow the standard convention of writing  $e(z)$  for  $e^{2\pi iz}$ ). Then on writing  $R(n)$  for the number of representations of  $n$  as the sum of  $s$   $k$ th powers of natural numbers, it follows from orthogonality that

$$R(n) = \int_0^1 f(\alpha)^s e(-\alpha n) d\alpha.$$

When  $\alpha$  is “close” to a rational number  $a/q$  with  $(a, q) = 1$  and  $q$  “small”, we expect that

$$f(\alpha) \sim q^{-1} S(q, a) v(\alpha - a/q),$$

where

$$S(q, a) = \sum_{r=1}^q e(ar^k/q) \quad \text{and} \quad v(\beta) = \int_0^P e(\beta \gamma^k) d\gamma.$$

This relation is straightforward to establish in an interval about  $a/q$ , so long as “close” and “small” are interpreted suitably. Now put

$$R_{\mathcal{A}}(n) = \int_{\mathcal{A}} f(\alpha)^s e(-\alpha n) d\alpha.$$

For a suitable union  $\mathfrak{M}$  of such intervals centred on  $a/q$  (the *major arcs*), and for  $s$  sufficiently large in terms of  $k$ , one can establish that as  $n \rightarrow \infty$ , the asymptotic relation

$$R_{\mathfrak{M}}(n) \sim \frac{\Gamma(1 + 1/k)^s}{\Gamma(s/k)} n^{s/k-1} \mathfrak{S}(n)$$

holds, where  $\mathfrak{S}(n)$  is the *singular series*

$$\mathfrak{S}(n) = \sum_{q=1}^{\infty} T(q; n)$$

and

$$T(q; n) = \sum_{\substack{a=1 \\ (a,q)=1}}^q q^{-s} S(q, a)^s e(-an/q).$$



## 8 The Necessary Congruence Condition

For each prime  $p$ , let

$$U(p; n) = \sum_{k=0}^{\infty} T(p^k; n).$$

The function  $T(q; n)$  is multiplicative. Thus, when the singular series converges absolutely, one has

$$\mathfrak{S}(n) = \prod_p U(p; n).$$

It is helpful to the success of the circle method that the singular series should satisfy  $\mathfrak{S}(n) \gg 1$ . With this observation in mind, Hardy and Littlewood [61] defined  $\Gamma(k)$  to be the least integer  $s$  with the property that, for every prime number  $p$ , there is a positive number  $C(p)$  such that  $U(p; n) \geq C(p)$  uniformly in  $n$ . Subsequently, Hardy and Littlewood [62] showed that indeed  $\mathfrak{S}(n) \gg 1$  whenever  $s \geq \max\{\Gamma(k), 4\}$ . Next let  $\Gamma_0(k)$  be the least number  $s$  with the property that the equation

$$x_1^k + \cdots + x_s^k = n \tag{8.1}$$

has a non-singular solution in  $\mathbb{Q}_p$  (or rather, that the corresponding congruence modulo  $q$  always has a solution with  $(x_1, q) = 1$ ). Hardy and Littlewood [63] were able to show that  $\Gamma_0(k) = \Gamma(k)$  (see Theorem 1 of the aforementioned paper). Thus one sees that the singular series reflects the local properties of sums of  $k$ -th powers. In particular, the singular series is zero whenever the equation (8.1) fails to have a  $p$ -adic solution, for some prime  $p$ , and this reflects the trivial observation that the equation can be soluble over  $\mathbb{Z}$  only if it is soluble everywhere locally.

Hardy and Littlewood [63] conjecture that  $\Gamma(k) \rightarrow \infty$  as  $k \rightarrow \infty$ , but it is not even known whether or not one has

$$\liminf_{k \rightarrow \infty} \Gamma(k) \geq 4.$$

When  $k > 2$ , they showed that  $\Gamma(k) = 4k$  when  $k$  is a power of 2 and that  $\Gamma(k) \leq 2k$  otherwise. They also computed  $\Gamma(k)$  exactly when  $3 \leq k \leq 36$ , and established that  $\Gamma(k) \geq 4$  when  $3 \leq k \leq 3000$ . Here they showed that equality occurs only when  $k = 3, 7, 19$ , and possibly (but improbably) when  $k = 1163, 1637, 1861, 1997, 2053$ . These values of  $k$  can probably be settled by modern computing methods, and doubtless the calculations could be carried a good deal further. As far as we are aware, nothing has been done in this direction.

For a more detailed exposition of the Hardy-Littlewood method and the analysis of the major arcs and the singular series, see Vaughan [128] (especially Chapters 2 and 4).

## 9 The Minor Arcs

In order to establish an asymptotic formula for  $R(n)$  it suffices to show that  $R_{\mathfrak{m}}(n) = o(n^{s/k-1})$ , where  $\mathfrak{m} = [0, 1) \setminus \mathfrak{M}$  (the *minor arcs*). One needs to show that the minor arc contribution  $R_{\mathfrak{m}}(n)$  is smaller by a factor  $o(n^{-1})$ , or equivalently  $o(P^{-k})$ , than the trivial estimate  $P^s$ . Routinely this is established via an inequality of the kind

$$\int_{\mathfrak{m}} |f(\alpha)|^s d\alpha \leq \left( \sup_{\mathfrak{m}} |f(\alpha)| \right)^{s-2t} \int_0^1 |f(\alpha)|^{2t} d\alpha.$$

The integral on the right hand side of this inequality may be interpreted as the number of solutions of an underlying diophantine equation, and it is from here that most of the savings usually come. On the other hand, non-trivial estimates for  $|f(\alpha)|$ , when  $\alpha \in \mathfrak{m}$ , may be obtained from estimates stemming from work of Weyl [154] and Vinogradov [144] (see Vaughan [128] for more modern estimates). When successful, this leads to the relation

$$R(n) \sim \frac{\Gamma(1 + 1/k)^s}{\Gamma(s/k)} n^{s/k-1} \mathfrak{S}(n). \quad (9.1)$$

It is in finding ways of dealing with the minor arcs, or in modifying the method so as to make the minor arcs more amenable, that most of the research has concentrated in the eighty years that have elapsed since the pioneering investigations of Hardy and Littlewood.

## 10 The Asymptotic Formula

Define  $\tilde{G}(k)$  to be the smallest natural number  $s_0$  such that whenever  $s \geq s_0$ , the asymptotic relation (9.1) holds. Work of Hardy and Littlewood [62], described already in §3, established the general upper bound

$$\tilde{G}(k) \leq (k-2)2^{k-1} + 5.$$

Progress on upper bounds for  $\tilde{G}(k)$  has since been achieved on two fronts. In Table 3 we present upper bounds for  $\tilde{G}(k)$  relevant for small values of  $k$ .

When  $k$  is large, bounds stemming from Vinogradov's mean value theorem provide dramatic improvements over the estimates recorded in Table 3.

Hua [75],	$2^k + 1,$
Vaughan [119][122],	$2^k \quad (k \geq 3),$
Heath-Brown [65][66],	$7 \cdot 2^{k-3} + 1 \quad (k \geq 6),$
Boklan [5],	$7 \cdot 2^{k-3} \quad (k \geq 6).$

**Table 3.** Upper bounds for  $\tilde{G}(k)$ : smaller  $k$

Vinogradov [142],	$183k^9(\log k + 1)^2 + 1,$
Vinogradov [141],	$91k^8(\log k + 1)^9 + 1 \quad (k \geq 20),$
Vinogradov [147],	$10k^2 \log k,$
Hua [77],	$4k^2(\log k + \frac{1}{2} \log \log k + 8),$
Wooley [156],	$2k^2(\log k + O(\log \log k)),$
Ford [53],	$k^2(\log k + \log \log k + O(1)).$

**Table 4.** Upper bounds for  $\tilde{G}(k)$ : larger  $k$

In Table 4 we present upper bounds for  $\tilde{G}(k)$  of use primarily when  $k$  is large.

We note that the methods underlying the last two bounds can be adapted to give explicit bounds for  $\tilde{G}(k)$  when  $k$  is of moderate size. Thus the method of Ford yields a bound for  $\tilde{G}(k)$  that is superior to the best recorded in Table 3 as soon as  $k \geq 9$ , and indeed unpublished work of Boklan and Wooley pushes this transition further to  $k \geq 8$ .

The bounds recorded in Tables 3 and 4 are likely to be a long way from the truth. One might expect that  $G(k) = \max\{k + 1, \Gamma_0(k)\}$ , and, with an appropriate interpretation of the asymptotic formula when  $\mathfrak{S}(n) = 0$ , that  $\tilde{G}(k) = k + 1$ .

One curiosity is that when  $k = 3$  and  $s = 7$ , it can be shown that  $R(n) \gg n^{4/3}$  (see Vaughan [125]), yet we are unable to show that  $R(n) \ll n^{4/3}$ . Indeed, it is currently the case that, quite generally, when  $s$  lies in the range between the known upper bounds for  $G(k)$  and  $\tilde{G}(k)$ , we can show that  $R(n) \gg n^{s/k-1}$ , but not  $R(n) \ll n^{s/k-1}$ .

## 11 Diminishing Ranges

In the Hardy-Littlewood method as outlined above the main problem is that of obtaining a suitable estimate for the mean value

$$\int_0^1 |f(\alpha)|^{2t} d\alpha,$$

that is, the number of integral solutions of the equation

$$x_1^k + \cdots + x_t^k = y_1^k + \cdots + y_t^k, \quad (11.1)$$

with  $1 \leq x_j, y_j \leq P$  ( $1 \leq j \leq t$ ). Available estimates can be improved significantly if the variables are restricted, an idea already present in Hardy and Littlewood [62]. Define

$$P_1 = \frac{1}{2}P, \quad P_j = \frac{1}{2}P_{j-1}^{1-1/k} \quad (2 \leq j \leq t),$$

and consider the equation (11.1) subject to the constraints  $P_j < x_j, y_j \leq 2P_j$  ( $1 \leq j \leq t$ ). Inspecting the expression  $|x_j^k - y_j^k|$  successively for  $j = 1, 2, \dots$  when  $x_j \neq y_j$ , we find that only the diagonal solutions in which  $x_j = y_j$  ( $1 \leq j \leq t$ ) occur. Thus we find that the number of solutions of this type is at most  $O(P_1 \dots P_t)$ . This saves  $P_1 \dots P_t$  over the trivial bound, which is of order  $(P_1 \dots P_t)^2$ . Now  $P_1 \dots P_t \approx P^\lambda$ , where

$$\lambda = 1 + (1 - 1/k) + \cdots + (1 - 1/k)^{t-1} = k - k(1 - 1/k)^t.$$

Already when  $t \sim Ck \log k$ , for a suitable positive constant  $C$ , this exponent is close to  $k$ . Vinogradov and Davenport have exploited and developed this idea in a number of ways (see, for example, Davenport [27][34], and Vinogradov [147][149]; see also Davenport and Erdős [35]).

There is a “ $p$ -adic” analogue of this idea, first exploited by Davenport [31], in which one considers expressions of the kind

$$x_1^k + p_2^k(x_2^k + p_3^k(x_3^k + \cdots))$$

on each side of the equation. Here the  $p_i$  denote suitably chosen prime numbers. The analysis rests on congruences of the type  $x_1 \equiv y_1 \pmod{p^k}$ . When  $p > P^{1/k}$  and  $1 \leq x_1, y_1 \leq P$ , this congruence implies that  $x_1 = y_1$ , and so on, just as in the diminishing ranges device. This idea has the merit of returning the various  $k$ -th powers  $p_2^k \dots p_j^k x_j^k$  to being in comparable size ranges. However in each of these methods the variables in (11.1) have varying natures and the homogeneity is essentially lost.

## 12 Smooth Numbers and Efficient Differences

In modern variants of the circle method as applied to Waring's problem, starting with the work of Vaughan [124], homogeneity is restored by considering the number of solutions  $S_t(P, R)$  of the equation (11.1) with  $x_j, y_j \in \mathcal{A}(P, R)$ , where  $\mathcal{A}(P, R)$  denotes the set of  $R$ -smooth numbers up to  $P$ , namely

$$\mathcal{A}(P, R) = \{n \in [1, P] \cap \mathbb{Z} : p|n \implies p \leq R\}.$$

In applications, one takes  $R$  to be a suitably small, but positive power of  $P$ . The set  $\mathcal{A}(P, R)$  has the extremely convenient property that, given any positive number  $M$  with  $M \leq P$ , and an element  $x \in \mathcal{A}(P, R)$  with  $x > M$ , there is always an integer  $m$  with  $m \in [M, MR]$  for which  $m|x$ . Moreover, this integer  $m$  can be coaxed into playing the rôle of the prime  $p$  in the  $p$ -adic argument mentioned above. Finally, and of great importance in what follows, the set  $\mathcal{A}(P, R)$  has positive density whenever  $R$  is no smaller than a positive power of  $P$ .

The objective now is to find good exponents  $\lambda_s$  with the property that whenever  $\varepsilon > 0$ , there exists a positive number  $\eta_0 = \eta_0(s, k, \varepsilon)$  such that whenever  $R = P^\eta$  with  $0 < \eta \leq \eta_0$ , one has

$$S_s(P, R) \ll P^{\lambda_s + \varepsilon}.$$

Such exponents are established via an iterative process in which a sequence of sets of exponents  $\boldsymbol{\lambda}^{(n)} = (\lambda_1^{(n)}, \lambda_2^{(n)}, \dots)$  is constructed by finding an expression for each  $\lambda_s^{(n+1)}$  in terms of the elements of  $\boldsymbol{\lambda}^{(n)}$ . Boundedness is trivial, so there is always a convergent subsequence. In fact, our arguments produce monotonicity, and the convergence is fairly rapid. For a more detailed introduction and motivation for the underlying ideas in using smooth numbers in Waring's problem, see the survey article Vaughan [127].

Beginning with the work of Wooley [155], a key element in the iterations is the repeated use of *efficient differencing*, and this procedure is fully exploited in subsequent work of Vaughan and Wooley [130] [131] [132] [134]. For each  $s \in \mathbb{N}$ , we take  $\phi_i = \phi_{i,s}$  ( $i = 1, \dots, k$ ) to be real numbers with  $0 \leq \phi_i \leq 1/k$ . For  $0 \leq j \leq k$ , we then define

$$P_j = 2^j P, \quad M_j = P^{\phi_j}, \quad H_j = P_j M_j^{-k}, \quad Q_j = P_j (M_1 \dots M_j)^{-1},$$

$$\tilde{H}_j = \prod_{i=1}^j H_i \quad \text{and} \quad \tilde{M}_j = \prod_{i=1}^j M_i R.$$

Define the modified forward difference operator,  $\Delta_j^*$ , recursively by taking

$$\Delta_1^*(f(x); h; m) = m^{-k}(f(x + hm^k) - f(x)),$$

and when  $j \geq 1$ , by inductively defining

$$\begin{aligned} \Delta_{j+1}^*(f(x); h_1, \dots, h_{j+1}; m_1, \dots, m_{j+1}) \\ = \Delta_1^*(\Delta_j^*(f(x); h_1, \dots, h_j; m_1, \dots, m_j); h_{j+1}; m_{j+1}). \end{aligned}$$

For  $0 \leq j \leq k$ , let

$$f(z) = (z - h_1 m_1^k - \dots - h_j m_j^k)^k,$$

and define the difference polynomial

$$\Psi_j = \Psi_j(z; h_1, \dots, h_j; m_1, \dots, m_j)$$

by taking

$$\Psi_j = \Delta_j^*(f(z); 2h_1, \dots, 2h_j; m_1, \dots, m_j).$$

Here we adopt the convention that  $\Psi_0(z; \mathbf{h}; \mathbf{m}) = z^k$ . We write

$$f_j(\alpha) = \sum_{x \in \mathcal{A}(Q_j, R)} e(\alpha x^k),$$

and

$$F_j(\alpha) = \sum_{z, \mathbf{h}, \mathbf{m}} e(\alpha \Psi_j(z; \mathbf{h}; \mathbf{m})),$$

where the summation is over  $z, \mathbf{h}, \mathbf{m}$  with

$$1 \leq z \leq P_j, \quad 1 \leq h_i \leq 2^{j-i} H_i,$$

$$M_i < m_i \leq M_i R, \quad m_i \in \mathcal{A}(P, R),$$

for  $1 \leq i \leq j$ . Finally, we define

$$T(j, s) = \int_0^1 |F_j(\alpha) f_j(\alpha)^{2s}| d\alpha.$$

Now, on considering the underlying diophantine equation, we have

$$S_{s+1}(P, R) \leq \int_0^1 |F_0(\alpha)^2 f_0(\alpha)^{2s}| d\alpha.$$

The starting point in the iterative process is to bound the latter expression in terms of  $S_s(Q_1, R)$  and  $T(1, s)$ . This corresponds to taking the first difference in the classical Weyl differencing argument, and extracting the contribution arising from those terms with  $x_1 = y_1$ . Thus one obtains

$$S_{s+1}(P, R) \ll P^\varepsilon M_1^{2s-1} (P M_1 S_s(Q_1, R) + T(1, s)),$$

and this inequality we write symbolically as

$$F_0^2 f_0^{2s} \mapsto F_1 f_1^{2s}.$$

One way to proceed is by means of a repeated efficient differencing step. In principle this is based on the Cauchy-Schwarz inequality, applied in the form

$$\int_0^1 |F_j f_j^{2s}| d\alpha \leq \left( \int_0^1 |f_j|^{2t} d\alpha \right)^{1/2} \left( \int_0^1 |F_j^2 f_j^{4s-2t}| d\alpha \right)^{1/2},$$

where for the sake of concision we have written  $f_j$  for  $f_j(\alpha)$  and likewise  $F_j$  for  $F_j(\alpha)$ . Thus, for  $j = 1, 2, \dots$ , the mean value  $T(j, s)$  can be related to  $S_t(Q_j, R)$  and  $T(j+1, 2s-t)$ , where  $t < 2s$  is a parameter at our disposal, via inequalities of the shape

$$T(j, s) \ll P^\varepsilon (S_t(Q_j, R))^{1/2} (\tilde{H}_j \tilde{M}_j M_{j+1}^{4s-2t-1} \Xi_{j+1})^{1/2},$$

where we write

$$\Xi_{j+1} = P \tilde{H}_j \tilde{M}_j M_{j+1} S_{2s-t}(Q_{j+1}, R) + T(j+1, 2s-t).$$

This is the  $(j+1)$ -th step in the differencing process and can be portrayed by

$$\begin{array}{ccc} F_j f_j^{2s} & \longrightarrow & F_{j+1} f_{j+1}^{4s-2t} \\ & & \downarrow \\ & & f_j^{2t} \end{array}$$

There are more sophisticated variants of this procedure wherein it may be useful to restrict some of the variables to a range  $(\frac{1}{2}Q_j R^{-j}, Q_j]$ , or to replace the set  $\mathcal{A}(Q_j, R)$  by  $\mathbb{N} \cap [1, Q_j]$  (see §2 of Vaughan and Wooley [134] for details, and a more complete discussion).

Another option is to use Hölder's inequality to bound  $T(j, s)$ . Thus we obtain an inequality of the type

$$T(j, s) = \int_0^1 |F_j(\alpha) f_j(\alpha)^{2s}| d\alpha \ll I_l^a I_{l+1}^b U_v^c U_w^d,$$

where

$$\begin{aligned} I_m &= \int_0^1 |F_j(\alpha)|^{2m} d\alpha \quad (m = l, l+1), \\ U_u &= \int_0^1 |f_j(\alpha)|^{2u} d\alpha \quad (u = v, w), \end{aligned}$$

and  $l, v, w, a, b, c, d$  are non-negative numbers satisfying the equations

$$a + b + c + d = 1, \quad 2^l a + 2^{l+1} b = 1, \quad vc + wd = s.$$

There is clearly great flexibility in the possible choices of the parameters here. We can summarise this process by

$$F_j f_j^{2s} \implies (F_j^{2^l})^a (F_j^{2^{l+1}})^b (f_j^{2v})^c (f_j^{2w})^d.$$

Yet another option is to apply the Hardy-Littlewood method to  $T(j, s)$ . In practice we expect that the minor arc contribution dominates, although this is not guaranteed. But if it does, then

$$T(j, s) \ll \left( \sup_{\alpha \in \mathfrak{m}} |F_j(\alpha)| \right) S_s(Q_j, R), \quad (12.1)$$

and this we abbreviate to

$$F_j f_j^{2s} \implies (F_j)(f_j^{2s}).$$

By optimising choices for the parameters in order to obtain the sharpest estimates at each stage of the iteration process, one ultimately obtains relations describing  $\lambda^{(n+1)}$  in terms of  $\lambda^{(n)}$ . The sharpest permissible exponents  $\lambda$  attainable by these methods are in general not easy to describe, and require substantial computations to establish (see, for example, Vaughan and Wooley [134]). However, one can describe in general terms the salient features of the permissible exponents  $\lambda_s$ . When  $s$  is rather small compared to  $k$ , it transpires that permissible exponents  $\lambda_s = s + \delta_s$  can be derived with  $\delta_s$  positive but small (see the next section for a consequence of this fact). Further, the simplest versions of the repeated efficient differencing method (see Wooley [155][157]) establish that the exponent  $\lambda_s = 2s - k + ke^{1-2s/k}$  is permissible for every natural number  $s$ . Roughly speaking, therefore, one may compare the respective strengths of the diminishing ranges argument, and the repeated efficient differencing method, by comparing how rapidly the respective functions  $k(1 - 1/k)^s \sim ke^{-s/k}$  and  $ke^{1-2s/k}$  tend to zero as  $s$  increases.

The improvements in the most recent work (Vaughan and Wooley [134]) come about mostly through the following technical improvements:

- Better use of the Hardy-Littlewood method to estimate

$$T(j, s) = \int_0^1 |F_j(\alpha) f_j(\alpha)^{2s}| d\alpha.$$

In particular, tighter control is exercised in mean on the behaviour of the exponential sum  $F_j(\alpha)$  on the major arcs, and this permits the assumption of (12.1) for a larger range of parameters  $\phi$  than previously available.



- Better estimates for

$$I_m = \int_0^1 |F_j(\alpha)|^{2m} d\alpha,$$

established largely via estimates for the number of integral points on certain affine plane curves.

- Better estimates (see Wooley [159]) for

$$\sup_{\alpha \in \mathfrak{m}} |f_0(\alpha)|.$$

Such estimates might be described as playing a significant rôle in the estimation of  $G(k)$  in the final stages of the analysis.

### 13 Breaking Classical Convexity

All of the methods thus far described depend, in a fundamental manner, on the natural interpretation of even moments of exponential sums in terms of the number of solutions of certain underlying diophantine equations. In §12, for example, one is limited to permissible exponents  $\lambda_s$  corresponding to integral values of  $s$ , and in this setting the most effective method for bounding odd and fractional moments of smooth Weyl sums is to apply Hölder's inequality to interpolate between even moments. With the natural extension of the notion of a permissible exponent  $\lambda_s$  from integral values of  $s$  to arbitrary positive numbers  $s$ , the resulting exponents form the convex hull of the set of permissible exponents  $\{\lambda_s : s \in \mathbb{N}\}$ . A perusal of §12 reveals that extra flexibility in choice of parameters, and therefore the potential for further improvements, will be achieved by the removal of this "classical convexity" barrier, and such has recently become available.

In Wooley [158], a method is established which, loosely speaking, enables one to replace the inequality

$$S_{s+1}(P, R) \ll P^\varepsilon M_1^{2s-1} (PM_1 S_s(Q_1, R) + T(1, s))$$

that occurred in §12 with  $s$  restricted to be a natural number, by the new inequality

$$S_{s+t}(P, R) \ll P^\varepsilon M_1^{2s-t} (P^t M_1^t S_s(Q_1, R) + T^*),$$

where

$$T^* = \int_0^1 |F(\alpha)^t f_1(\alpha)^{2s}| d\alpha$$

and

$$F(\alpha) = \sum_{\substack{u \in \mathcal{A}(M_1 R, R) \\ u > M_1}} \sum_{\substack{z_1, z_2 \in \mathcal{A}(P, R) \\ z_1 \equiv z_2 \pmod{u^k} \\ z_1 \neq z_2}} e(\alpha u^{-k}(z_1^k - z_2^k)).$$

In this latter estimate, the parameter  $s$  is no longer restricted to be integral, and the parameter  $t$  may be chosen freely with  $0 < t \leq 1$ . Moreover, the mean value  $T^*$  is very much reminiscent of  $T(1, s)$ , with  $F(\alpha)$  substituted for  $F_1(\alpha)$  and exhibiting similar properties. Thus, in addition to removing the integrality constraint on  $s$ , one may also iterate with a fractional number  $2t$  of variables.

As might be expected, the additional flexibility gained in this way leads to improved permissible exponents  $\lambda_s$  even for integral  $s$ , since our methods are so highly iterative. The overall improvements are usually quite small and are largest for smaller values of  $k$ . Such progress has not yet delivered sharper bounds for  $G(k)$ , but this work provides the sharpest results available concerning sums of cubes (see §4, and also Brüdern and Wooley [18]), and has also permitted new conclusions to be derived in certain problems involving sums of mixed powers (see Brüdern and Wooley [16] [19] [17], and also §15 below). Also, this “breaking convexity” device provides the best available lower bounds for the number  $\mathcal{N}_{k,s}(X)$  of natural numbers not exceeding  $X$  that are the sum of  $s$   $k$ th powers of positive integers, at least when  $s$  is small compared to  $k$ . Thus, when  $2 < s \leq 2e^{-1}\sqrt{k}$ , one has

$$\mathcal{N}_{k,s}(X) \gg X^{s/k - e^{-\gamma_s k}},$$

where  $\gamma_s = 16/(es)^2$  (see Theorems 1.3 and 1.4 of Wooley [158], and the associated discussion). For comparison, one conjecturally has the lower bound  $\mathcal{N}_{k,s}(X) \gg X^{s/k}$  whenever  $s < k$ .

Much remains to be investigated for fractional moments, in part owing to the substantial increase in complexity of the underlying computations (see Wooley [161] for more on this). However, such developments presently appear unlikely to have a large impact on the central problem of bounding  $G(k)$  in the classical version of Waring’s problem.

## 14 Variants of Waring’s Problem: Primes

Much work has been devoted to various generalisations of the classical version of Waring’s problem, and it seems appropriate to discuss some of the more mainstream variants.

We begin with the Waring-Goldbach problem, in which one seeks to represent integers as sums of  $k$ th powers of prime numbers. In order to

describe the associated local conditions, suppose that  $k$  is a natural number and  $p$  is a prime number. We denote by  $\theta = \theta(k, p)$  the integer with  $p^\theta | k$  and  $p^{\theta+1} \nmid k$ , and then define  $\gamma = \gamma(k, p)$  by

$$\gamma(k, p) = \begin{cases} \theta + 2, & \text{when } p = 2 \text{ and } \theta > 0, \\ \theta + 1, & \text{otherwise.} \end{cases}$$

Finally, we put

$$K(k) = \prod_{(p-1)|k} p^\gamma.$$

Denote by  $H(k)$  the least integer  $s$  such that every sufficiently large positive integer congruent to  $s$  modulo  $K(k)$  may be written as a sum of  $s$   $k$ th powers of prime numbers. Note that when  $(p-1)|k$ , one has  $p^\theta(p-1)|k$ , whence  $a^k \equiv 1 \pmod{p^\gamma}$  whenever  $(p, a) = 1$ . This explains our seemingly awkward definition of  $H(k)$ , since whenever  $n$  is the sum of  $s$   $k$ th powers of primes exceeding  $k+1$ , then necessarily  $n \equiv s \pmod{K(k)}$ . Naturally, further congruence conditions may arise from primes  $p$  with  $(p-1) \nmid k$ .

Following the pioneering investigations of Vinogradov [145][146] (see also Vinogradov [147]), Hua comprehensively investigated additive problems involving prime numbers in his influential book (see Hua [79], but also Hua [78]). Thus, it is known that for every natural number  $k$  one has

$$H(k) \leq 2^k + 1,$$

and, when  $k$  is large, that

$$H(k) \leq 4k(\log k + \frac{1}{2} \log \log k + O(1)).$$

In the conventional plan of attack on the Waring-Goldbach problem, one applies the Hardy-Littlewood method in a manner similar to that outlined above, but in interpreting the number of solutions of an analogue of the equation (11.1) over prime numbers, one obtains an upper bound by discarding the primality condition. With sufficiently many variables employed to save a factor of  $n$  via such an approach, one additional variable suffices to save the extra power of  $\log n$  required by primality considerations. Although this strategy evidently prohibits the use of smooth numbers, the diminishing ranges technology perfected by Davenport, and refined by Vaughan [121] and Thanigasalam [109]–[113] plays a prominent rôle in establishing the best available upper bounds for  $H(k)$  when  $k$  is small. We should also mention that recent progress depends on good estimates of Weyl-type for exponential sums over primes, and allied sums, available from the use of Vaughan's identity (see Vaughan [117]), combined

with the linear sieve equipped with a switching principle (see Kawada and Wooley [83]). Thus, for  $4 \leq k \leq 10$ , the best known upper bounds for  $H(k)$  are as follows:

$$\begin{aligned} \text{Kawada and Wooley [83], } & H(4) \leq 14, H(5) \leq 21, \\ \text{Thanigasalam [111], } & H(6) \leq 33, H(7) \leq 47, H(8) \leq 63, \\ & H(9) \leq 83, H(10) \leq 107. \end{aligned}$$

Despite much effort on the Waring-Goldbach problem for exponents 1, 2 and 3, further progress remains elusive. Improvements are feasible, however, if one is prepared to accept almost-primes in place of prime numbers (see, in particular, Chen [25], Brüdern [13][14], and Brüdern and Fouvry [15]). Difficulties related to those associated with the Waring-Goldbach problem are encountered when other sequences are substituted for prime numbers. For Waring's problem with smooth variables, see Harcos [57] and Brüdern and Wooley [18]. Also, see Nechaev [97] for work on Waring's problem with polynomial summands (Wooley [160] and Ford [55] have restricted improvements employing smooth numbers).

## 15 Variants of Waring's Problem: Sums of Mixed Powers

Suppose that  $k_1, k_2, \dots, k_s$  are natural numbers with  $2 \leq k_1 \leq k_2 \leq \dots \leq k_t$ . Then an optimistic counting argument suggests that whenever the equation

$$x_1^{k_1} + x_2^{k_2} + \dots + x_s^{k_s} = n \tag{15.1}$$

has  $p$ -adic solutions for each prime  $p$ , and

$$k_1^{-1} + k_2^{-1} + \dots + k_s^{-1} > 1, \tag{15.2}$$

then  $n$  should be represented as the sum of mixed powers of positive integers (15.1) whenever it is sufficiently large in terms of  $\mathbf{k}$ . When  $s = 3$  such an assertion may fail in certain circumstances (see Jagy and Kaplansky [80], or Exercise 5 of Chapter 8 of Vaughan [128]), but a heuristic application of the Hardy-Littlewood method suggests, at least, that the condition (15.2) should ensure that *almost all* integers in the expected congruence classes are thus represented. Moreover, subject instead to the condition

$$k_1^{-1} + k_2^{-1} + \dots + k_s^{-1} > 2, \tag{15.3}$$

a formal application of the circle method suggests that *all* integers in the expected congruence classes should be represented in the form (15.1). Meanwhile, a simple counting argument shows that in circumstances in which

the condition (15.2) does not hold, then arbitrarily large integers are not represented in the desired form.

The investigation of such analogues of Waring's problem for mixed powers has, since the early days of the Hardy-Littlewood method, stimulated progress in technology of use even in the classical version of Waring's problem. Additive problems in which the summands are restricted to be squares, cubes or biquadrates are perhaps of greater interest than those with higher powers, and here the current situation is remarkably satisfactory. We summarise below the current state of knowledge in the simpler problems of this nature. In Tables 5 and 6 we list constellations of powers whose sum represents, respectively, almost all, and all, integers subject to the expected congruence conditions. The tables are arranged, roughly speaking, starting with predominantly smaller exponents, and ending with predominantly larger exponents, and therefore not in chronological order of the results.

We have been unable to trace the origin in the literature of the conclusion on a square, two biquadrates and a  $k$ th power, but refer the reader to Exercise 6 of §2.8 of Vaughan [128] for related ideas (see also Roth

Davenport & Heilbronn [36],	two squares, one $k$ th power,
Davenport & Heilbronn [37],	one square, two cubes,
Roth [101],	one square, one cube, one biquadrate,
Vaughan [116],	one square, one cube, one fifth power,
Folklore (?),	one square, two biquadrates, one $k$ th power,
Hooley [70],	one square, one cube, one sixth power, one $k$ th power,
Davenport [29],	four cubes,
Brüdern [9][8],	three cubes, one biquadrate,
Brüdern [8], Lu [89],	three cubes, one fifth power,
Brüdern & Wooley [19],	three cubes, one sixth power ,
Kawada & Wooley [82],	one cube, four biquadrates,
Vaughan [124],	six biquadrates,
Kawada & Wooley [82],	five biquadrates, one $k$ th power ( $k$ odd).

**Table 5.** Representation of almost all integers

Gauss [56],	three squares,
Hooley [71],	two squares, three cubes,
Hooley [70],	two squares, assorted powers,
Vaughan [123],	one square, five cubes,
Brüdern & Wooley [16],	one square, four cubes, one biquadrate,
Ford [54],	one square, one cube, one biquadrate, ...,
	one fifteenth power,
Linnik [87][88],	seven cubes,
Brüdern [9],	six cubes, two biquadrates,
Brüdern [9],	five cubes, three biquadrates,
Kawada & Wooley [82],	three cubes, six biquadrates,
Brüdern & Wooley [17],	two cubes, seven biquadrates,
Kawada & Wooley [82],	one cube, nine biquadrates
Vaughan [124],	twelve biquadrates,
Kawada & Wooley [82],	ten biquadrates, one $k$ th power
	( $k$ odd).

**Table 6.** Representation of all integers

[101]). We remark that all ternary problems of interest have been solved, since for non-trivial triples  $(k_1, k_2, k_3)$  not accounted for in Table 5, one has  $k_1^{-1} + k_2^{-1} + k_3^{-1} \leq 1$ . Also, energetic readers may be interested in tackling a problem which presently defies resolution only by the narrowest of margins, namely the problem of showing that almost all integers are represented as the sum of two cubes and two biquadrates of positive integers.

Note that although the three square theorem is commonly ascribed to Legendre, his “proof” depended on an unsubstantiated assumption only later established by Dirichlet, and the first complete proof is due to Gauss. We finish by noting that in problems involving sums of two squares, methods more effective than the circle method can be brought into play (see especially Hooley [70][71] and Brüdern [7]).

## 16 Variants of Waring’s Problem: Beyond $\mathbb{Z}$

Given the considerable energy expended on the investigation of Waring’s problem over the rational integers, it seems natural to extend this work

to algebraic number fields. Here one encounters the immediate difficulty of deciding what precisely Waring's problem should mean in this broader context. It is possible, for example, that an algebraic integer in a number field  $K$  may not be a sum of any finite number of  $k$ th powers of algebraic integers of that field (consider, say, the parity of the imaginary part of  $\mathfrak{a}^2$  when  $\mathfrak{a} \in \mathbb{Z}[i]$ ). With this in mind, when  $K$  is a number field and  $\mathfrak{O}_K$  is its ring of integers, we define  $J_k$  to be the subring of  $\mathfrak{O}_K$  generated by the  $k$ th powers of integers of  $K$ . We must also provide an analogue of the positivity of the  $k$ th powers inherent in the classical version of Waring's problem. Thus we define  $G_K(k)$  to be the smallest positive integer  $s$  with the property that, for some positive number  $c = c(k, K)$ , and for all totally positive integers  $\nu \in J_k$  of sufficiently large norm, the equation

$$\nu = \lambda_1^k + \lambda_2^k + \cdots + \lambda_s^k \quad (16.1)$$

is always soluble in totally non-negative integers  $\lambda_j$  of  $K$  with  $N(\lambda_j) \leq cN(\nu)^{1/k}$  ( $1 \leq j \leq s$ ).

Following early work of Meissner [93] and Mordell [95] for a restricted class of number fields, Siegel [102][103] was the first to obtain quite general conclusions for sums of squares, and hence, via the method of Hilbert [69], for sums of  $k$ th powers. Siegel later developed a proper generalisation of the Hardy-Littlewood method to number fields, and here the dissection into major and minor arcs is a particular source of difficulty. In this way, Siegel [104][105] obtained the upper bound

$$G_K(k) \leq dk(2^{k-1} + d) + 1,$$

where  $d$  denotes the degree of the field  $K$ . If one were to break the equation (16.1) into components with respect to an integral basis for  $\mathfrak{O}_K$ , then one would obtain  $d$  equations of degree  $k$  in  $ds$  variables, and so one might optimistically expect the analytic part of the circle method to apply with a number of variables roughly the same in both  $K$  and  $\mathbb{Q}$ . Perhaps motivated by such considerations, Siegel asked for reasonable bounds on  $G_K(k)$  independent of  $d$ . This question was ultimately addressed through the work of Birch [4] and Ramanujam [98], who provided the upper bound

$$G_K(k) \leq \max\{8k^5, 2^k + 1\}. \quad (16.2)$$

It is evident that the uniform bound (16.2) is far from the above cited bound  $G(k) \leq (1 + o(1))k \log k$  of Wooley [155], and the only slightly weaker precursors of Vinogradov. An important desideratum, therefore, is the reduction of the bound (16.2) to one of similar order to that presently available for  $G(k)$ , or at least a reduction to a bound polynomial in  $k$ .

Failing this, effort has been expended in pursuit of bounds of order  $k \log k$ , but with a modest dependence on the degree  $d$  of the field  $K$ . Progress towards this objective has mirrored developments in the classical version of Waring's problem. Thus, building on work generalising that of Vinogradov to the number field setting by Körner [84] and Eda [50], methods employing smooth numbers and repeated efficient differences have recently been applied by Davidson [39] to establish the bound

$$G_K(k) \leq (3 + o(1))k \log k + c_d k,$$

where

$$c_d = 4d + 3 \log\left(\frac{1}{2}(d^2 + 1)\right) + 7,$$

and the term  $o(1)$  is independent of  $d$  (apparently, the term  $3 + o(1)$  here can be replaced by  $2 + o(1)$  with only modest effort). Moreover, when  $K$  has class number one, Davidson [40][42] has obtained the bound

$$G_K(k) \leq k(\log k + \log \log k + c_d),$$

where  $c_d$  is approximately  $4d$ . Finally, Davidson [41] has improved on earlier work of Tatzuza [106] to establish the strikingly simple conclusion that, for every number field  $K$ , one has

$$G_K(k) \leq 2d(\widehat{G}(k) + 2k),$$

where  $\widehat{G}(k)$  denotes the least number  $s$  satisfying the property that the set of rational integers that can be expressed as the sum of  $s$   $k$ th powers of natural numbers has positive density (in particular, of course,  $\widehat{G}(k) \leq G_1(k)$ ).

More exotic still than the variants of Waring's problem over number fields are those in which one works over the polynomial rings  $\mathbb{F}_q[t]$ . Here one must again impose restrictions on the size of polynomials employed in the representation (and in this situation, the degree of a polynomial provides a measure of its size). Analogues of the Hardy-Littlewood method have been devised in this polynomial ring setting (see, for example, Effinger and Hayes [51]). Unfortunately, however, Weyl differencing proves ineffective whenever  $k$ th powers are considered over  $\mathbb{F}_q[t]$  with  $k \geq \text{char}(\mathbb{F}_q)$ , for in such circumstances the factor  $k!$ , introduced into the argument of the exponential sum over  $k$ th powers via the differencing argument, is equal to zero in  $\mathbb{F}_q[t]$ . Consequently, one frequently restricts attention to  $k$ th powers with  $k$  smaller than the characteristic (but see Car and Cherly [22] for results on sums of 11 cubes in  $\mathbb{F}_{2^h}[t]$ ). With this restriction, Car [20][21] has shown that every polynomial  $M$ , with  $M \in \mathbb{F}_q[t]$ , of sufficiently large degree, can be written in the form

$$M = M_1^k + \cdots + M_s^k,$$



with  $M_i \in \mathbb{F}_q[t]$  of degree smaller than  $1 + (\deg(M))/k$  ( $1 \leq s \leq k$ ), provided that

$$s \geq \min\{2^k + 1, 2k(k-1) \log 2 + 2k + 3\}.$$

## 17 Open Problems and Conjectures

Returning temporarily to the methods of §12, there are a number of problems connected with the mean value

$$I_m = \int_0^1 |F_j(\alpha)|^{2m} d\alpha$$

which suggest some interesting questions. We will concentrate on the situation with  $m = 1$ , and remark only that the available results become less satisfactory as  $m$  increases.

A simple combinatorial argument reveals that the difference polynomial  $\Psi_j = \Psi_j(z; \mathbf{h}; \mathbf{m})$  is given explicitly by the formula

$$\Psi_j = k!2^j h_1 \dots h_j \sum_{u \geq 0} \sum_{v_1 \geq 0} \dots \sum_{v_j \geq 0} \frac{z^u (h_1 m_1^k)^{2v_1} \dots (h_j m_j^k)^{2v_j}}{u!(2v_1 + 1)! \dots (2v_j + 1)!},$$

where the summation is subject to the condition  $u + 2v_1 + \dots + 2v_j = k - j$ . Consequently, one has

$$\Psi_j = h_1 \dots h_j z^d \sum_{r=0}^{\frac{1}{2}(k-j-d)} c_r(h_1 m_1^k, \dots, h_j m_j^k) z^{2r},$$

where

$$d = \begin{cases} 0, & \text{when } k - j \text{ is even,} \\ 1, & \text{when } k - j \text{ is odd,} \end{cases}$$

and for  $0 \leq r \leq (k - j - d)/2$ , the coefficients  $c_r(\boldsymbol{\xi})$  are polynomials with positive integral coefficients that are symmetric in  $\xi_1^2, \dots, \xi_j^2$ , and have total degree  $k - j - 2r - d$ . Now the mean value

$$I_1 = \int_0^1 |F_j(\alpha)|^2 d\alpha$$

is equal to the number of solutions of the diophantine equation

$$\Psi_j(z; \mathbf{h}; \mathbf{m}) = \Psi_j(z'; \mathbf{h}'; \mathbf{m}'), \quad (17.1)$$

with the variables in ranges discernible from the definition of  $F_j(\alpha)$  in §12, and one might hope that the total number of solutions is close to the number of diagonal solutions, which is to say that

$$I_1 \ll P^{1+\varepsilon} \widetilde{M}_j \widetilde{H}_j.$$

When  $k-j$  is odd (so  $d = 1$ ), the presence of the term  $z^d$  makes it especially easy to deal with equation (17.1) by exploiting the inherent multiplicative structure, and indeed one can achieve the desired bound provided also that  $j \leq (k-d)/3$ . The cases  $j = 1$  and  $k-j = 2$  or  $4$  are also doable. However, when  $k-j$  is even and  $1 < j \leq k-6$ , the situation is not so easy. By the way, this difficulty already occurs in Davenport's work [31]. To illustrate this situation, the simplest special case that we cannot handle directly corresponds to  $k = 8$  and  $j = 2$ , and here one has

$$\begin{aligned} \Psi_j &= h_1 h_2 (224z^6 + 1120z^4 (h_1^2 m_1^{16} + h_2^2 m_2^{16}) \\ &\quad + z^2 (672h_1^4 m_1^{32} + 2240h_1^2 h_2^2 m_1^{16} m_2^{16} + 672h_2^4 m_2^{32}) \\ &\quad + 32h_1^6 m_1^{48} + 224h_1^4 h_2^2 m_1^{32} m_2^{16} + 224h_1^2 h_2^4 m_1^{16} m_2^{32} + 32h_2^6 m_2^{48}). \end{aligned}$$

This suggests various general questions.

- Suppose that  $f, g \in \mathbb{Z}[x]$ . Are there simple conditions on  $f, g$  such that the number  $N$  of integral points  $(x, y) \in [-P, P]^2$  for which  $f(x) = g(y)$  satisfies

$$N \ll (P\mathcal{H}(f)\mathcal{H}(g))^\varepsilon?$$

Here  $\mathcal{H}(h)$  denotes the height of  $h$ . A qualitative version of this has already been considered by Davenport, Lewis and Schinzel [38], and if  $f(x) - g(y)$  is irreducible over  $\mathbb{C}$ , then a celebrated theorem of Siegel shows that the number of solutions is finite unless there is a rational parametric solution of special form.

By the way, in view of the above, it is perhaps not surprising that in our treatment of  $I_m$  the bound of Bombieri and Pila [6] plays a rôle.

- Suppose that  $\mathcal{A} \subset \mathbb{Z}^k \cap [-X, X]^k$ . Let  $R(n; \mathcal{A})$  denote the number of solutions of the equation

$$a_1 x + a_2 x^2 + \cdots + a_k x^k = n$$

with  $x \in \mathbb{Z} \cap [-P, P]$  and  $\mathbf{a} \in \mathcal{A}$ . Are there any simple conditions under which it is true that

$$\sum_n R(n; \mathcal{A})^2 \ll \text{card}(\mathcal{A}) P(XP)^\varepsilon?$$

• A well-known conjecture in connection with Waring's problem is Hypothesis K (Hardy and Littlewood [62]). Let

$$R_{k,s}(n) = \text{card}\{\mathbf{x} \in \mathbb{N}^s : x_1^k + \cdots + x_s^k = n\}.$$

Then Hypothesis K asserts that for each natural number  $k$ , one has

$$R_{k,k}(n) \ll n^\varepsilon. \quad (17.2)$$

From this it would follow that

$$G(k) \leq \max\{2k + 1, \Gamma_0(k)\} \quad (17.3)$$

and

$$G_1(k) = \max\{k + 1, \Gamma_0(k)\}. \quad (17.4)$$

The conjecture (17.2) was later shown by Mahler [90] to be false for  $k = 3$ , and indeed his counter-example shows that, infinitely often, one has  $R_{3,3}(n) > 9^{-1/3}n^{1/12}$ . However, the conjecture is still open when  $k \geq 4$ , and for (17.3) and (17.4), it suffices to know that

$$\sum_{n \leq N} R_{k,k}(n)^2 \ll N^{1+\varepsilon}.$$

Hooley [74] established this when  $k = 3$ , under the assumption of the Riemann Hypothesis for a certain Hasse-Weil  $L$ -function. As far as we know, no simple conjecture of this kind is known from which it would follow that  $G(k) = \max\{k + 1, \Gamma_0(k)\}$ .

• It may well be true that, when  $k \geq 3$ , one has

$$\sum_{n \leq N} R_{k,k}(n)^2 \sim CN.$$

However Hooley [72] has shown, at least when  $k = 3$ , that the constant  $C$  here is larger than what would arise simply from the major arcs. This leads to some interesting speculations. The number of solutions of the equation

$$x_1^k + \cdots + x_s^k = y_1^k + \cdots + y_s^k \leq N,$$

in which the variables on the right hand side are a permutation of those on the left hand side, is asymptotic to  $C_1 N^{s/k}$ , for a certain positive number  $C_1 = C_1(k, s)$ , and when  $s < k$  the contribution arising from the major arcs is smaller. Maybe one should think of these solutions as being "trivial", "parametric", or as arising from some "degenerate" property of the geometry of the surface. Anyway, their contribution is mostly concentrated

on the minor arcs. It seems rather likely that this phenomenon persists for  $s \geq k$  and explains Hooley's discovery. This leads to the philosophy that the major arcs correspond to non-trivial solutions, and the minor arcs to trivial solutions. There is an example of this phenomenon in Vaughan and Wooley [133].

- Recall the definitions of  $f(\alpha)$ ,  $S(q, a)$  and  $v(\beta)$  from §7. One can conjecture that, whenever  $(a, q) = 1$ , one has

$$f(\alpha) - q^{-1}S(q, a)v(\alpha - a/q) \ll (q + P^k|q\alpha - a|)^{1/k}. \quad (17.5)$$

Possibly the exponent has to be weakened to  $1/k + \varepsilon$ , but any counterexamples would be interesting.

From (17.5) it would follow that

$$\sum_{n \leq N} R_{k,k}(n)^2 \ll N.$$

Also it is just conceivable that (17.5), in combination with a variant of the Hardy-Littlewood-Kloosterman method, would achieve the bound  $G(k) \leq \max\{\mathfrak{G}(k), \Gamma_0(k)\}$ , where  $\mathfrak{G}(k) < 2k + 1$ .

- The inequality (17.5) is a special case of conjectures that can be made about the exponential sum

$$f(\alpha) = \sum_{x \leq P} e(\alpha_1 x + \cdots + \alpha_k x^k)$$

that would have many consequences in analytic number theory. For example one can ask if something like

$$f(\alpha) - q^{-1}S(q, \mathbf{a})v(\alpha - \mathbf{a}/q) \ll \sum_{j=2}^k \left( \frac{q + P^j |\alpha_j q - a_j|}{(q, a_j)} \right)^{1/j} \quad (17.6)$$

is true. Here

$$S(q, \mathbf{a}) = \sum_{r=1}^q e((a_1 r + \cdots + a_k r^k)/q),$$

$$v(\beta) = \int_0^P e(\beta_1 \gamma + \cdots + \beta_k \gamma^k) d\gamma,$$

and  $a_1 = c(q, a_2, \dots, a_k)$ , where  $c = c(q, \mathbf{a})$  is the unique integer with

$$-\frac{1}{2} < c - \alpha_1 q / (q, a_2, \dots, a_k) \leq \frac{1}{2}.$$

The inequality (17.6) is known to hold for  $k = 2$  (Vaughan unpublished) with the right hand side weakened slightly to

$$\left(\frac{q}{(q, a_2)}\right)^{1/2} \left(\log \frac{2q}{(q, a_2)} + P|\alpha_2 - a_2/q_2|^{1/2}\right).$$

• One way of viewing the Hardy-Littlewood method is that we begin by considering the Fourier transform with respect to Lebesgue measure on  $[0, 1)$  for an appropriate generating function defined in terms of the additive characters, and then approximate to it by a product of discrete measures. Since part of the problem when one has relatively few variables is that the geometry genuinely intrudes, one can ask whether we are using the best measure for the problem at hand. In this situation something more closely related to the underlying geometry might be more useful.

In conclusion, it is clear that although we have come a long way in the twentieth century, there remains plenty still to be done!

## References

- [1] M. A. Bennett, *Fractional parts of powers of rational numbers*, Math. Proc. Cambridge Philos. Soc. **114** (1993), 191–201.
- [2] ———, *An ideal Waring problem with restricted summands*, Acta Arith. **66** (1994), 125–132.
- [3] F. Beukers, *Fractional parts of powers of rationals*, Math. Proc. Cambridge Philos. Soc. **90** (1981), 13–20.
- [4] B. J. Birch, *Waring's problem in algebraic number fields*, Proc. Cambridge Philos. Soc. **57** (1961), 449–459.
- [5] K. D. Boklan, *The asymptotic formula in Waring's problem*, Matematika **41** (1994), 147–161.
- [6] E. Bombieri and J. Pila, *The number of integral points on arcs and ovals*, Duke Math. J. **59** (1989), 337–357.
- [7] J. Brüdern, *Sums of squares and higher powers*, J. London Math. Soc. (2) **35** (1987), 233–243.
- [8] ———, *Iterationsmethoden in der additiven Zahlentheorie*, Dissertation, Göttingen, 1988.

- [9] ———, *On Waring's problem for cubes and biquadrates*, J. London Math. Soc. (2) **37** (1988), 25–42.
- [10] ———, *Sums of four cubes*, Monatsh. Math. **107** (1989), 179–188.
- [11] ———, *On Waring's problem for fifth powers and some related topics*, Proc. London Math. Soc. (3) **61** (1990), 457–479.
- [12] ———, *On Waring's problem for cubes*, Math. Proc. Cambridge Philos. Soc. **109** (1991), 229–256.
- [13] ———, *Sieves, the circle method, and Waring's problem for cubes*, Habilitationsschrift, Göttingen, 1991, Mathematica Göttingensis **51**.
- [14] ———, *A sieve approach to the Waring-Goldbach problem, II. On the seven cube theorem*, Acta Arith. **72** (1995), 211–227.
- [15] J. Brüdern and É. Fouvry, *Lagrange's four squares theorem with almost prime variables*, J. Reine Angew. Math. **454** (1994), 59–96.
- [16] J. Brüdern and T. D. Wooley, *On Waring's problem: a square, four cubes and a biquadrate*, Math. Proc. Cambridge Philos. Soc. **127** (1999), 193–200.
- [17] ———, *On Waring's problem: two cubes and seven biquadrates*, Tsukuba J. Math. **24** (2000), 387–417.
- [18] ———, *On Waring's problem for cubes and smooth Weyl sums*, Proc. London Math. Soc. (3) **82** (2001), 89–109.
- [19] ———, *On Waring's problem: three cubes and a sixth power*, Nagoya Math. J. **163** (2001), 13–53.
- [20] M. Car, *Le problème de Waring pour l'anneau des polynômes sur un corps fini*, Séminaire de Théorie des Nombres 1972–1973 (Univ. Bordeaux I, Talence), Lab. Théorie des Nombres, CNRS, Talence, 1973, Exp. No. 6, 13pp.
- [21] ———, *Waring's problem in function fields*, Proc. London Math. Soc. (3) **68** (1994), 1–30.
- [22] M. Car and J. Cherly, *Sommes de cubes dans l'anneau  $\mathbb{F}_2^h[X]$* , Acta Arith. **65** (1993), 227–241.
- [23] J. W. S. Cassels and R. C. Vaughan, *Ivan Matveevich Vinogradov*, Biographical Memoirs of Fellows of the Royal Society **31** (1985), 613–631, reprinted in: Bull. London Math. Soc. **17** (1985), 584–600.

- [24] J.-R. Chen, *On Waring's problem for  $n$ -th powers*, Acta Math. Sinica **8** (1958), 253–257.
- [25] ———, *On the representation of a large even integer as the sum of a prime and the product of at most two primes*, Sci. Sinica **16** (1973), 157–176.
- [26] R. J. Cook, *A note on Waring's problem*, Bull. London Math. Soc. **5** (1973), 11–12.
- [27] H. Davenport, *Sur les sommes de puissances entières*, C. R. Acad. Sci. Paris, Sér. A **207** (1938), 1366–1368.
- [28] ———, *On sums of positive integral  $k$ th powers*, Proc. Roy. Soc. London, Ser. A **170** (1939), 293–299.
- [29] ———, *On Waring's problem for cubes*, Acta Math. **71** (1939), 123–143.
- [30] ———, *On Waring's problem for fourth powers*, Ann. of Math. **40** (1939), 731–747.
- [31] ———, *On sums of positive integral  $k$ th powers*, Amer. J. Math. **64** (1942), 189–198.
- [32] ———, *On Waring's problem for fifth and sixth powers*, Amer. J. Math. **64** (1942), 199–207.
- [33] ———, *Sums of three positive cubes*, J. London Math. Soc. **25** (1950), 339–343.
- [34] ———, *The collected works of Harold Davenport, Volume III (B. J. Birch, H. Halberstam and C. A. Rogers, eds.)*, Academic Press, London, 1977.
- [35] H. Davenport and P. Erdős, *On sums of positive integral  $k$ th powers*, Ann. of Math. **40** (1939), 533–536.
- [36] H. Davenport and H. Heilbronn, *Note on a result in the additive theory of numbers*, Proc. London Math. Soc. (2) **43** (1937), 142–151.
- [37] ———, *On Waring's problem: two cubes and one square*, Proc. London Math. Soc. (2) **43** (1937), 73–104.
- [38] H. Davenport, D. J. Lewis, and A. Schinzel, *Equations of the form  $f(x) = g(y)$* , Quart. J. Math. Oxford (2) **12** (1961), 304–312.

- [39] M. Davidson, *Sums of  $k$ -th powers in number fields*, *Mathematika* **45** (1998), 359–370.
- [40] ———, *A new minor-arcs estimate for number fields*, *Topics in number theory* (University Park, PA, 1997), Kluwer Acad. Publ., Dordrecht, 1999, pp. 151–161.
- [41] ———, *On Waring’s problem in number fields*, *J. London Math. Soc.* (2) **59** (1999), 435–447.
- [42] ———, *On Siegel’s conjecture in Waring’s problem*, to appear, 2001.
- [43] J.-M. Deshouillers, F. Hennecart, K. Kawada, B. Landreau, and T. D. Wooley, *Sums of biquadrates: a survey*, in preparation, 2001.
- [44] J.-M. Deshouillers, F. Hennecart, and B. Landreau, 7, 373, 170, 279, 850, *Math. Comp.* **69** (2000), 421–439.
- [45] ———, *Waring’s problem for sixteen biquadrates – Numerical results*, *J. Théorie Nombres Bordeaux* **12** (2000), 411–422, *Colloque International de Théorie des Nombres* (Talence, 1999).
- [46] J.-M. Deshouillers, K. Kawada, and T. D. Wooley, *On sums of sixteen biquadrates*, in preparation, 2001.
- [47] L. E. Dickson, *All integers except 23 and 239 are sums of eight cubes*, *Bull. Amer. Math. Soc.* **45** (1939), 588–591.
- [48] ———, *History of the theory of numbers, Vol. II: Diophantine analysis*, Chelsea, New York, 1966.
- [49] A. K. Dubitskas, *A lower bound on the value  $\|(3/2)^k\|$* , *Uspekhi Mat. Nauk* **45** (1990), 153–154, translation in: *Russian Math. Surveys* **45** (1990), 163–164.
- [50] Y. Eda, *On Waring’s problem in an algebraic number field*, *Rev. Columbiana Math.* **9** (1975), 29–73.
- [51] G. Effinger and D. Hayes, *Additive number theory of polynomials over a finite field*, Oxford University Press, Oxford, 1991.
- [52] T. Estermann, *On Waring’s problem for fourth and higher powers*, *Acta Arith.* **2** (1937), 197–211.
- [53] K. B. Ford, *New estimates for mean values of Weyl sums*, *Internat. Math. Res. Notices* (1995), 155–171.



- [54] ———, *The representation of numbers as sums of unlike powers. II*, J. Amer. Math. Soc. **9** (1996), 919–940.
- [55] ———, *Waring's problem with polynomial summands*, J. London Math. Soc. (2) **61** (2000), 671–680.
- [56] C. F. Gauss, *Disquisitiones arithmeticae*, Leipzig, 1801.
- [57] G. Harcos, *Waring's problem with small prime factors*, Acta Arith. **80** (1997), 165–185.
- [58] G. H. Hardy and J. E. Littlewood, *A new solution of Waring's problem*, Quart. J. Math. Oxford **48** (1920), 272–293.
- [59] ———, *Some problems of "Partitio Numerorum": I. A new solution of Waring's problem*, Göttingen Nachrichten (1920), 33–54.
- [60] ———, *Some problems of "Partitio Numerorum": II. Proof that every large number is the sum of at most 21 biquadrates*, Math. Z. **9** (1921), 14–27.
- [61] ———, *Some problems of "Partitio Numerorum": IV. The singular series in Waring's Problem and the value of the number  $G(k)$* , Math. Z. **12** (1922), 161–188.
- [62] ———, *Some problems of "Partitio Numerorum" (VI): Further researches in Waring's problem*, Math. Z. **23** (1925), 1–37.
- [63] ———, *Some problems of "Partitio Numerorum" (VIII): The number  $\Gamma(k)$  in Waring's problem*, Proc. London Math. Soc. (2) **28** (1928), 518–542.
- [64] G. H. Hardy and S. Ramanujan, *Asymptotic formulae in combinatory analysis*, Proc. London Math. Soc. (2) **17** (1918), 75–115.
- [65] D. R. Heath-Brown, *Weyl's inequality, Hua's inequality, and Waring's problem*, J. London Math. Soc. (2) **38** (1988), 396–414.
- [66] ———, *Weyl's inequality and Hua's inequality*, Number Theory (Ulm, 1987), Lecture Notes in Math., vol. 1380, Springer-Verlag, Berlin, 1989, pp. 87–92.
- [67] ———, *The circle method and diagonal cubic forms*, Philos. Trans. Roy. Soc. London Ser. A **356** (1998), 673–699.
- [68] H. Heilbronn, *Über das Waringsche Problem*, Acta Arith. **1** (1936), 212–221.

- [69] D. Hilbert, *Beweis für Darstellbarkeit der ganzen Zahlen durch eine feste Anzahl nter Potenzen Waringsche Problem*, Nach. Königl. Ges. Wiss. Göttingen math.-phys. Kl. (1909), 17–36, Math. Ann. **67** (1909), 281–300.
- [70] C. Hooley, *On a new approach to various problems of Waring's type*, Recent progress in analytic number theory, Vol. 1 (Durham, 1979), Academic Press, London, 1981, pp. 127–191.
- [71] ———, *On Waring's problem for two squares and three cubes*, J. Reine Angew. Math. **328** (1981), 161–207.
- [72] ———, *On some topics connected with Waring's problem*, J. Reine Angew. Math. **369** (1986), 110–153.
- [73] ———, *On Waring's problem*, Acta Math. **157** (1986), 49–97.
- [74] ———, *On Hypothesis  $K^*$  in Waring's problem*, Sieve methods, exponential sums, and their applications in number theory (Cardiff, 1995), Cambridge Univ. Press, Cambridge, 1997, pp. 175–185.
- [75] L.-K. Hua, *On Waring's problem*, Quart. J. Math. Oxford **9** (1938), 199–202.
- [76] ———, *Some results on Waring's problem for smaller powers*, C. R. Acad. Sci. URSS (2) **18** (1938), 527–528.
- [77] ———, *An improvement of Vinogradov's mean-value theorem and several applications*, Quart. J. Math. Oxford **20** (1949), 48–61.
- [78] ———, *Die Abschätzungen von Exponentialsummen und ihre Anwendung in der Zahlentheorie*, B.G. Teubner, Leipzig, 1959.
- [79] ———, *Additive theory of prime numbers*, American Math. Soc., Providence, Rhode Island, 1965.
- [80] W. C. Jagy and I. Kaplansky, *Sums of squares, cubes, and higher powers*, Experiment. Math. **4** (1995), 169–173.
- [81] R. D. James, *On Waring's problem for odd powers*, Proc. London Math. Soc. **37** (1934), 257–291.
- [82] K. Kawada and T. D. Wooley, *Sums of fourth powers and related topics*, J. Reine Angew. Math. **512** (1999), 173–223.
- [83] ———, *On the Waring-Goldbach problem for fourth and fifth powers*, Proc. London Math. Soc. (3) **83** (2001), 1–50.

- [84] O. Körner, *Über das Waringsche Problem in algebraischen Zahlkörpern*, Math. Ann. **144** (1961), 224–238.
- [85] J. M. Kubina and M. C. Wunderlich, *Extending Waring's conjecture to 471, 600, 000*, Math. Comp. **55** (1990), 815–820.
- [86] E. Landau, *Über eine Anwendung der Primzahlen auf das Waringsche Problem in der elementaren Zahlentheorie*, Math. Ann. **66** (1909), 102–105.
- [87] Ju. V. Linnik, *On the representation of large numbers as sums of seven cubes*, Dokl. Akad. Nauk SSSR **35** (1942), 162.
- [88] ———, *On the representation of large numbers as sums of seven cubes*, Mat. Sb. **12** (1943), 218–224.
- [89] M. G. Lu, *On Waring's problem for cubes and fifth power*, Sci. China Ser. A **36** (1993), 641–662.
- [90] K. Mahler, *Note on hypothesis K of Hardy and Littlewood*, J. London Math. Soc. **11** (1936), 136–138.
- [91] ———, *On the fractional parts of the powers of a rational number (II)*, Mathematika **4** (1957), 122–124.
- [92] K. S. McCurley, *An effective seven cube theorem*, J. Number Theory **19** (1984), 176–183.
- [93] O. E. Meissner, *Über die Darstellung der Zahlen einiger algebraischen Zahlkörper als Summen von Quadratzahlen des Körpers*, Arch. Math. Phys. (3) **7** (1904), 266–268.
- [94] Z. Z. Meng, *Some new results on Waring's problem*, J. China Univ. Sci. Tech. **27** (1997), 1–5.
- [95] L. J. Mordell, *On the representation of algebraic numbers as a sum of four squares*, Proc. Cambridge Philos. Soc. **20** (1921), 250–256.
- [96] V. Narasimhamurti, *On Waring's problem for 8th, 9th, and 10th powers*, J. Indian Math. Soc. (N.S.) **5** (1941), 122.
- [97] V. I. Nechaev, *Waring's problem for polynomials*, Trudy Mat. Inst. Steklov, Izdat. Akad. Nauk SSSR, Moscow **38** (1951), 190–243.
- [98] C. P. Ramanujam, *Sums of  $m$ -th powers in  $p$ -adic rings*, Mathematika **10** (1963), 137–146.

- [99] K. Sambasiva Rao, *On Waring's problem for smaller powers*, J. Indian Math. Soc. **5** (1941), 117–121.
- [100] C. J. Ringrose, *Sums of three cubes*, J. London Math. Soc. (2) **33** (1986), 407–413.
- [101] K. F. Roth, *Proof that almost all positive integers are sums of a square, a positive cube and a fourth power*, J. London Math. Soc. **24** (1949), 4–13.
- [102] C. L. Siegel, *Darstellung total positiver Zahlen durch Quadrate*, Math. Z. **11** (1921), 246–275.
- [103] ———, *Additive Theorie der Zahlkörper II*, Math. Ann. **88** (1923), 184–210.
- [104] ———, *Generalisation of Waring's problem to algebraic number fields*, Amer. J. Math. **66** (1944), 122–136.
- [105] ———, *Sums of  $m$ -th powers of algebraic integers*, Ann. of Math. **46** (1945), 313–339.
- [106] T. Tatzuzawa, *On Waring's problem in algebraic number fields*, Acta Arith. **24** (1973), 37–60.
- [107] K. Thanigasalam, *On Waring's problem*, Acta Arith. **38** (1980), 141–155.
- [108] ———, *Some new estimates for  $G(k)$  in Waring's problem*, Acta Arith. **42** (1982), 73–78.
- [109] ———, *Improvement on Davenport's iterative method and new results in additive number theory, I*, Acta Arith. **46** (1985), 1–31.
- [110] ———, *Improvement on Davenport's iterative method and new results in additive number theory, II. Proof that  $G(5) \leq 22$* , Acta Arith. **46** (1986), 91–112.
- [111] ———, *Improvement on Davenport's iterative method and new results in additive number theory, III*, Acta Arith. **48** (1987), 97–116.
- [112] ———, *On sums of positive integral powers and simple proof of  $G(6) \leq 31$* , Bull. Calcutta Math. Soc. **81** (1989), 279–294.
- [113] ———, *On admissible exponents for  $k$ th powers*, Bull. Calcutta Math. Soc. **86** (1994), 175–178.

- [114] K.-C. Tong, *On Waring's problem*, *Advancement in Math.* **3** (1957), 602–607.
- [115] R. C. Vaughan, *Homogeneous additive equations and Waring's problem*, *Acta Arith.* **33** (1977), 231–253.
- [116] ———, *A ternary additive problem*, *Proc. London Math. Soc.* (3) **41** (1980), 516–532.
- [117] ———, *Recent work in additive prime number theory*, *Proceedings of the International Congress of Mathematicians (Helsinki, 1978)*, *Acad. Sci. Fennica*, 1980, pp. 389–394.
- [118] ———, *Sums of three cubes*, *Bull. London Math. Soc.* **17** (1985), 17–20.
- [119] ———, *On Waring's problem for cubes*, *J. Reine Angew. Math.* **365** (1986), 122–170.
- [120] ———, *On Waring's problem for sixth powers*, *J. London Math. Soc.* (2) **33** (1986), 227–236.
- [121] ———, *On Waring's problem for smaller exponents*, *Proc. London Math. Soc.* (3) **52** (1986), 445–463.
- [122] ———, *On Waring's problem for smaller exponents. II*, *Mathematika* **33** (1986), 6–22.
- [123] ———, *On Waring's problem: one square and five cubes*, *Quart. J. Math. Oxford* (2) **37** (1986), 117–127.
- [124] ———, *A new iterative method in Waring's problem*, *Acta Math.* **162** (1989), 1–71.
- [125] ———, *On Waring's problem for cubes II*, *J. London Math. Soc.* (2) **39** (1989), 205–218.
- [126] ———, *A new iterative method in Waring's problem II*, *J. London Math. Soc.* (2) **39** (1989), 219–230.
- [127] ———, *The use in additive number theory of numbers without large prime factors*, *Philos. Trans. Roy. Soc. London Ser. A* **345** (1993), 363–376.
- [128] ———, *The Hardy-Littlewood Method, 2nd edition*, *Cambridge University Press*, Cambridge, 1997.

- [129] R. C. Vaughan and T. D. Wooley, *On Waring's problem: some refinements*, Proc. London Math. Soc. (3) **63** (1991), 35–68.
- [130] ———, *Further improvements in Waring's problem, III: Eighth powers*, Philos. Trans. Roy. Soc. London Ser. A **345** (1993), 385–396.
- [131] ———, *Further improvements in Waring's problem, II: Sixth powers*, Duke Math. J. **76** (1994), 683–710.
- [132] ———, *Further improvements in Waring's problem*, Acta Math. **174** (1995), 147–240.
- [133] ———, *On a certain nonary cubic form and related equations*, Duke Math. J. **80** (1995), 669–735.
- [134] ———, *Further improvements in Waring's problem, IV: Higher powers*, Acta Arith. **94** (2000), 203–285.
- [135] I. M. Vinogradov, *A new estimate for  $G(n)$  in Waring's problem*, Dokl. Akad. Nauk SSSR **5** (1934), 249–253.
- [136] ———, *A new solution of Waring's problem*, Dokl. Akad. Nauk SSSR **2** (1934), 337–341.
- [137] ———, *On some new results in the analytic theory of numbers*, C. R. Acad. Sci. Paris, Sér. A **199** (1934), 174–175.
- [138] ———, *On the upper bound for  $G(n)$  in Waring's problem*, Izv. Akad. Nauk SSSR Ser. Fiz.-Matem. **10** (1934), 1455–1469.
- [139] ———, *A new variant of the proof of Waring's theorem*, Trud. Matem. instituta it. V. A. Steklov **9** (1935), 5–15.
- [140] ———, *A new variant of the proof of Waring's theorem*, C. R. Acad. Sci. Paris, Sér. A **200** (1935), 182–184.
- [141] ———, *An asymptotic formula for the number of representations in Waring's problem*, Mat. Sb. **42** (1935), 531–534.
- [142] ———, *New estimates for Weyl sums*, Dokl. Akad. Nauk SSSR **8** (1935), 195–198.
- [143] ———, *On Waring's problem*, Ann. of Math. **36** (1935), 395–405.
- [144] ———, *On Weyl's sums*, Mat. Sb. **42** (1935), 521–530.
- [145] ———, *Representation of an odd number as a sum of three primes*, Dokl. Akad. Nauk SSSR **15** (1937), 6–7.

- [146] ———, *Some theorems concerning the theory of primes*, Mat. Sb. **44** (1937), 179–195.
- [147] ———, *The method of trigonometrical sums in the theory of numbers*, Trav. Inst. Math. Stekloff **23** (1947), 109 pp.
- [148] ———, *On an upper bound for  $G(n)$* , Izv. Akad. Nauk SSSR Ser. Mat. **23** (1959), 637–642.
- [149] ———, *Selected Works (translated from the Russian by N. Psv)*, Springer-Verlag, Berlin, 1985.
- [150] E. Waring, *Meditationes Algebraicæ, second edition*, Archdeacon, Cambridge, 1770.
- [151] ———, *Meditationes Algebraicæ, third edition*, Archdeacon, Cambridge, 1782.
- [152] ———, *Meditationes Algebraicæ*, American Math. Soc., Providence, 1991, translation by Dennis Weeks of the 1782 edition.
- [153] G. L. Watson, *A proof of the seven cube theorem*, J. London Math. Soc. **26** (1951), 153–156.
- [154] H. Weyl, *Über die Gleichverteilung von Zahlen mod Eins*, Math. Ann. **77** (1916), 313–352.
- [155] T. D. Wooley, *Large improvements in Waring's problem*, Ann. of Math. **135** (1992), 131–164.
- [156] ———, *On Vinogradov's mean value theorem*, Mathematika **39** (1992), 379–399.
- [157] ———, *The application of a new mean value theorem to the fractional parts of polynomials*, Acta Arith. **65** (1993), 163–179.
- [158] ———, *Breaking classical convexity in Waring's problem: sums of cubes and quasi-diagonal behaviour*, Invent. Math. **122** (1995), 421–451.
- [159] ———, *New estimates for smooth Weyl sums*, J. London Math. Soc. (2) **51** (1995), 1–13.
- [160] ———, *On exponential sums over smooth numbers*, J. Reine Angew. Math. **488** (1997), 79–140.
- [161] ———, *Quasi-diagonal behaviour and smooth Weyl sums*, Monatsh. Math. **130** (2000), 161–170.

[162] ———, *Sums of three cubes*, *Mathematika*, to appear, 2001.