Jacobi triple product and the like via the exclusion process and the like

Joint with Ross Bowen, Dan Fretwell, Jessica Jay

Márton Balázs

University of Bristol

Interacting Particle Systems and Hydrodynamic Limits CRM 23 March, 2022.

Jacobi triple product

Theorem Let |x| < 1 and $y \neq 0$ be complex numbers. Then

$$\prod_{i=1}^{\infty} (1-x^{2i}) \left(1+\frac{x^{2i-1}}{y^2}\right) \left(1+x^{2i-1}y^2\right) = \sum_{m=-\infty}^{\infty} x^{m^2} y^{2m}.$$

Mostly appears in number theory and combinatorics of partitions.

We'll prove it using interacting particles (for real x, y only).

Models

Asymmetric simple exclusion Zero range

Blocking measures

State space No boundaries Boundaries

Lay down - stand up

Jacobi triple product

More models





Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.


Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.



Particles try to jump

to the right with rate p, to the left with rate q = 1 - p < p.





Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump


Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump



Particles jump

Product blocking measures

Can we have a reversible stationary distribution in product form:

$$\underline{\mu}(\underline{\omega}) = \bigotimes_{i} \mu_{i}(\omega_{i});$$

$$\underline{\mu}(\underline{\omega}) \cdot \mathsf{rate}(\underline{\omega} \to \underline{\omega}^{i \frown i+1}) = \underline{\mu}(\underline{\omega}^{i \frown i+1}) \cdot \mathsf{rate}(\underline{\omega}^{i \frown i+1} \to \underline{\omega}) \quad ?$$

Here

$$\underline{\omega}^{i \frown i+1} = \underline{\omega} - \underline{\delta}_i + \underline{\delta}_{i+1}.$$

Asymmetric simple exclusion



Asymmetric simple exclusion



$$\underline{\mu}(\underline{\omega}) \cdot \mathsf{rate}(\underline{\omega} \to \underline{\omega}^{i \frown i+1}) = \underline{\mu}(\underline{\omega}^{i \frown i+1}) \cdot \mathsf{rate}(\underline{\omega}^{i \frown i+1} \to \underline{\omega}) \quad ?$$

AZRP:

 $\mu_i(\omega_i)\mu_{i+1}(\omega_{i+1})\cdot p\mathbf{1}\{\omega_i>0\}=\mu_i(\omega_i-1)\mu_{i+1}(\omega_{i+1}+1)\cdot q$

Solution:
$$\mu_i \sim \text{Geometric}\Big(1 - \Big(\frac{p}{q}\Big)^{i-\text{const}}\Big).$$

Notice:

$$\mathbf{P}\{\eta_i = 0\} = 1 - \varrho_i = \frac{1}{1 + (\frac{\rho}{q})^{i-c}} \qquad \text{as } i \to \infty$$
$$\mathbf{P}\{\eta_i = 1\} = \varrho_i = \frac{1}{(\frac{q}{p})^{i-c} + 1} \qquad \text{as } i \to -\infty$$

are both summable. Hence by Borel-Cantelli there is μ -a.s. a rightmost hole and a leftmost particle,

$$N := \sum_{i=1}^{\infty} (1 - \eta_i) - \sum_{i=-\infty}^{0} \eta_i$$

is μ -a.s. finite.



Also: it is conserved. Define

$$\Omega^n := \{\mathbf{0}, \mathbf{1}\}^{\mathbb{Z}} \cap \{\mathbf{N}(\underline{\eta}) = \mathbf{n}\},\$$

$$\underline{\mu}\Big(\bigcup_{n=-\infty}^{\infty}\Omega^n\Big)=1$$



Also: it is conserved. Define

$$\Omega^n := \{\mathbf{0}, \mathbf{1}\}^{\mathbb{Z}} \cap \{\mathbf{N}(\underline{\eta}) = \mathbf{n}\},\$$

$$\underline{\mu}\Big(\bigcup_{n=-\infty}^{\infty}\Omega^n\Big)=1$$



Also: it is conserved. Define

$$\Omega^n := \{\mathbf{0}, \mathbf{1}\}^{\mathbb{Z}} \cap \{\mathbf{N}(\underline{\eta}) = \mathbf{n}\},\$$

$$\underline{\mu}\Big(\bigcup_{n=-\infty}^{\infty}\Omega^n\Big)=1$$



Also: it is conserved. Define

$$\Omega^n := \{\mathbf{0}, \mathbf{1}\}^{\mathbb{Z}} \cap \{\mathbf{N}(\underline{\eta}) = \mathbf{n}\},\$$

$$\underline{\mu}\Big(\bigcup_{n=-\infty}^{\infty}\Omega^n\Big)=1$$



Also: it is conserved. Define

$$\Omega^n := \{\mathbf{0}, \mathbf{1}\}^{\mathbb{Z}} \cap \{\mathbf{N}(\underline{\eta}) = \mathbf{n}\},\$$

$$\underline{\mu}\Big(\bigcup_{n=-\infty}^{\infty}\Omega^n\Big)=1$$



Also: it is conserved. Define

$$\Omega^n := \{\mathbf{0}, \mathbf{1}\}^{\mathbb{Z}} \cap \{\mathbf{N}(\underline{\eta}) = \mathbf{n}\},\$$

$$\underline{\mu}\Big(\bigcup_{n=-\infty}^{\infty}\Omega^n\Big)=1$$



Also: it is conserved. Define

$$\Omega^n := \{\mathbf{0}, \mathbf{1}\}^{\mathbb{Z}} \cap \{\mathbf{N}(\underline{\eta}) = \mathbf{n}\},\$$

$$\underline{\mu}\Big(\bigcup_{n=-\infty}^{\infty}\Omega^n\Big)=1$$



Also: it is conserved. Define

$$\Omega^n := \{\mathbf{0}, \mathbf{1}\}^{\mathbb{Z}} \cap \{\mathbf{N}(\underline{\eta}) = \mathbf{n}\},\$$

$$\underline{\mu}\Big(\bigcup_{n=-\infty}^{\infty}\Omega^n\Big)=1$$



Also: it is conserved. Define

$$\Omega^n := \{\mathbf{0}, \mathbf{1}\}^{\mathbb{Z}} \cap \{\mathbf{N}(\underline{\eta}) = \mathbf{n}\},\$$

$$\underline{\mu}\Big(\bigcup_{n=-\infty}^{\infty}\Omega^n\Big)=1$$



Also: it is conserved. Define

$$\Omega^n := \{\mathbf{0}, \mathbf{1}\}^{\mathbb{Z}} \cap \{\mathbf{N}(\underline{\eta}) = \mathbf{n}\},\$$

$$\underline{\mu}\Big(\bigcup_{n=-\infty}^{\infty}\Omega^n\Big)=1$$



Also: it is conserved. Define

$$\Omega^n := \{\mathbf{0}, \mathbf{1}\}^{\mathbb{Z}} \cap \{\mathbf{N}(\underline{\eta}) = \mathbf{n}\},\$$

$$\underline{\mu}\Big(\bigcup_{n=-\infty}^{\infty}\Omega^n\Big)=1$$

Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.


Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



 $\tau : \Omega^n \to \Omega^{n-1}$

Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



 $\tau : \Omega^n \to \Omega^{n-1}$

Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



 $\tau : \Omega^n \to \Omega^{n-1}$

Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



 $\tau : \Omega^n \to \Omega^{n-1}$

Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



 $\tau : \Omega^n \to \Omega^{n-1}$

Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



 $\tau : \Omega^n \to \Omega^{n-1}$

Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



 $\tau : \Omega^n \to \Omega^{n-1}$

Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



 $\tau : \Omega^n \to \Omega^{n-1}$

Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



 $\tau : \Omega^n \to \Omega^{n-1}$

Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.



 $\tau : \Omega^n \to \Omega^{n-1}$

Left shift:
$$(\tau \underline{\eta})_i = \eta_{i+1}$$
.





Ergodic decomposition of $\underline{\mu}$ with $\underline{\nu}^{n}(\cdot) := \underline{\mu}(\cdot | \mathbf{N}(\cdot) = n)$.

Let's find the coefficients $\underline{\mu}(N(\cdot) = n)!$

$$\varrho_i = \frac{(\frac{p}{q})^{i-c}}{1 + (\frac{p}{q})^{i-c}} = \frac{1}{(\frac{q}{p})^{i-c} + 1}$$

$$\varrho_i = \frac{\left(\frac{p}{q}\right)^{i-c}}{1 + \left(\frac{p}{q}\right)^{i-c}} = \frac{1}{\left(\frac{q}{p}\right)^{i-c} + 1}$$
$$\underline{\mu}(\underline{\eta}) = \prod_{i \le 0} \frac{\left(\frac{p}{q}\right)^{(i-c)\eta_i}}{1 + \left(\frac{p}{q}\right)^{i-c}} \cdot \prod_{i>0} \frac{\left(\frac{q}{p}\right)^{(i-c)(1-\eta_i)}}{\left(\frac{q}{p}\right)^{i-c} + 1}$$

$$\varrho_{i} = \frac{\left(\frac{p}{q}\right)^{i-c}}{1 + \left(\frac{p}{q}\right)^{i-c}} = \frac{1}{\left(\frac{q}{p}\right)^{i-c} + 1}$$
$$\underline{\mu}(\underline{\eta}) = \prod_{i \le 0} \frac{\left(\frac{p}{q}\right)^{(i-c)\eta_{i}}}{1 + \left(\frac{p}{q}\right)^{i-c}} \cdot \prod_{i > 0} \frac{\left(\frac{q}{p}\right)^{(i-c)(1-\eta_{i})}}{\left(\frac{q}{p}\right)^{i-c} + 1}$$
$$\underline{\mu}(\underline{\tau}\underline{\eta}) = \frac{\prod_{i \le 0} \left(\frac{p}{q}\right)^{(i-c)\eta_{i+1}}}{\prod_{i \le 0} \left(1 + \left(\frac{p}{q}\right)^{i-c}\right)} \cdot \frac{\prod_{i > 0} \left(\frac{q}{p}\right)^{(i-c)(1-\eta_{i+1})}}{\prod_{i > 0} \left(\left(\frac{q}{p}\right)^{i-c} + 1\right)}$$

$$\varrho_{i} = \frac{\left(\frac{p}{q}\right)^{i-c}}{1+\left(\frac{p}{q}\right)^{i-c}} = \frac{1}{\left(\frac{q}{p}\right)^{i-c}+1}$$

$$\underline{\mu}(\underline{\eta}) = \prod_{i \leq 0} \frac{\left(\frac{p}{q}\right)^{(i-c)\eta_{i}}}{1+\left(\frac{p}{q}\right)^{i-c}} \cdot \prod_{i>0} \frac{\left(\frac{q}{p}\right)^{(i-c)(1-\eta_{i})}}{\left(\frac{q}{p}\right)^{i-c}+1}$$

$$\underline{\mu}(\underline{\tau}\underline{\eta}) = \frac{\prod_{i \leq 0} \left(\frac{p}{q}\right)^{(i-c)\eta_{i+1}}}{\prod_{i \leq 0} \left(1+\left(\frac{p}{q}\right)^{i-c}\right)} \cdot \frac{\prod_{i>0} \left(\frac{q}{p}\right)^{(i-c)(1-\eta_{i+1})}}{\prod_{i>0} \left(\left(\frac{q}{p}\right)^{i-c}+1\right)}$$

$$= \dots = \underline{\mu}(\underline{\eta}) \cdot \left(\frac{p}{q}\right)^{N(\underline{\eta})-c}.$$

$$\underline{\mu}(N=n-1)=\sum_{\underline{\eta}:N(\underline{\eta})=n-1}\underline{\mu}(\underline{\eta})$$

$$\underline{\mu}(N = n - 1) = \sum_{\underline{\eta}: N(\underline{\eta}) = n - 1} \underline{\mu}(\underline{\eta})$$
$$= \sum_{\underline{\eta}: N(\underline{\tau}\underline{\eta}) = n - 1} \underline{\mu}(\underline{\tau}\underline{\eta})$$

$$\underline{\mu}(N = n - 1) = \sum_{\underline{\eta}: N(\underline{\eta}) = n - 1} \underline{\mu}(\underline{\eta})$$
$$= \sum_{\underline{\eta}: N(\underline{\tau}\underline{\eta}) = n - 1} \underline{\mu}(\underline{\tau}\underline{\eta})$$
$$= \sum_{\eta: N(\eta) = n} \underline{\mu}(\underline{\eta}) \cdot \left(\frac{p}{q}\right)^{n - c}$$

$$\underline{\mu}(N = n - 1) = \sum_{\underline{\eta}: N(\underline{\eta}) = n - 1} \underline{\mu}(\underline{\eta})$$
$$= \sum_{\underline{\eta}: N(\underline{\tau}\underline{\eta}) = n - 1} \underline{\mu}(\underline{\tau}\underline{\eta})$$
$$= \sum_{\eta: N(\eta) = n} \underline{\mu}(\underline{\eta}) \cdot \left(\frac{p}{q}\right)^{n - c}$$

$$\underline{\mu}(N = n - 1) = \sum_{\underline{\eta}: N(\underline{\eta}) = n - 1} \underline{\mu}(\underline{\eta})$$
$$= \sum_{\underline{\eta}: N(\underline{\eta}) = n - 1} \underline{\mu}(\underline{\tau}\underline{\eta})$$
$$= \sum_{\underline{\eta}: N(\underline{\eta}) = n} \underline{\mu}(\underline{\eta}) \cdot \left(\frac{\underline{p}}{q}\right)^{n - c}$$
$$= \underline{\mu}(N = n) \cdot \left(\frac{\underline{p}}{q}\right)^{n - c}.$$

So,

$$\underline{\mu}(N = n - 1) = \sum_{\underline{\eta}: N(\underline{\eta}) = n - 1} \underline{\mu}(\underline{\eta})$$
$$= \sum_{\underline{\eta}: N(\underline{\eta}) = n - 1} \underline{\mu}(\underline{\tau}\underline{\eta})$$
$$= \sum_{\underline{\eta}: N(\underline{\eta}) = n} \underline{\mu}(\underline{\eta}) \cdot \left(\frac{\underline{p}}{q}\right)^{n - c}$$
$$= \underline{\mu}(N = n) \cdot \left(\frac{\underline{p}}{q}\right)^{n - c}.$$

Solution:

$$\underline{\mu}(N=n) = \frac{\left(\frac{q}{p}\right)^{\frac{n^2+n}{2}-cn}}{\sum \left(\frac{q}{p}\right)^{\frac{m^2+m}{2}-cm}}$$

discrete Gaussian.

and, if $N(\underline{\eta}) = n$,

$$\underline{\nu}^{n}(\underline{\eta}) = \underline{\mu}(\underline{\eta} \mid N(\underline{\eta}) = n) = \frac{\underline{\mu}(\underline{\eta})}{\mu(N(\underline{\eta}) = n)}$$
$$= \frac{\prod_{i \leq 0} \left(\frac{p}{q}\right)^{(i-c)\eta_{i}}}{\prod_{i \leq 0} \left(1 + \left(\frac{p}{q}\right)^{i-c}\right)} \cdot \frac{\prod_{i>0} \left(\frac{q}{p}\right)^{(i-c)(1-\eta_{i})}}{\prod_{i>0} \left(\left(\frac{q}{p}\right)^{i-c} + 1\right)} \cdot \frac{\sum_{i>0} \left(\frac{q}{p}\right)^{\frac{m^{2}+m}{2}-cm}}{\left(\frac{q}{p}\right)^{\frac{m^{2}+m}{2}-cn}}.$$

This is the unique stationary distribution on Ω^n .

Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\left(1 - \left(rac{p}{q}
ight)^{i-\operatorname{const}}
ight).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\left(1 - \left(rac{p}{q}
ight)^{i-\operatorname{const}}
ight).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$


Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \text{Geometric} \Big(1 - \Big(rac{p}{q} \Big)^{i - \text{const}} \Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$



Recall: Stationary distribution with marginals

$$\mu_i \sim \operatorname{Geometric}\Big(1 - \Big(rac{p}{q}\Big)^{i-\operatorname{const}}\Big).$$

 \rightarrow we have a problem: cannot do this for all *i*! We'll pick const = 1 and have a *right boundary* instead.



~> The product measure stays stationary on the half-line.

Lay down / stand up

ASEP



Lay down / stand up

ASEP





Lay down / stand up

ASEP




















ASEP



-2 -1 0 i









































































since stationary distributions of countable irreducible Markov chains are unique.

Jacobi triple product















































































































 \oplus to the right





 \oplus to the right





 \oplus to the right





 \oplus to the right





 \oplus to the right





 \oplus to the right





 \oplus to the right





 \oplus to the right





 \oplus to the right





 \oplus to the right





 \oplus to the right













































 \ominus to the left





 \ominus to the left





 \ominus to the left





 \ominus to the left





 \ominus to the left





 \ominus to the left





 \ominus to the left





 \ominus to the left





 \ominus to the left





 \ominus to the left














$\oplus \ominus \emptyset$ models





























$\oplus \ominus \emptyset$ models





... and the same steps to the left.





























No two consecutive 0's!

Odd ground state:



Even ground state:



No two consecutive 0's! is a nonlocal constraint.

- ▶ *No two consecutive 0's!* is a nonlocal constraint.
- The stood up model is nice otherwise. It has reversible product blocking measures.

- ▶ No two consecutive 0's! is a nonlocal constraint.
- The stood up model is nice otherwise. It has reversible product blocking measures.
- Reversible measures survive forbidden jumps. Yay!

The identities we get have to do with *generalised Frobenius partitions* and *generalised Young diagrams*.

ASEP(q, 1) Carinci, Giardiná, Redig, Sasamoto:

Jump	Rate	Jump	Rate
$\oplus \rightarrow$	$q^{-1} + q^{-3}$	$\leftarrow \oplus$	$q+q^3$
$\leftarrow \ominus$	$q^{-1} + q^{-3}$	$\ominus \rightarrow$	$q+q^3$
$\emptyset \leadsto \ominus \oplus$	q^{-3}	$\emptyset \leadsto \oplus \ominus$	q^3
$\oplus \ominus \leadsto \emptyset$	$(1+q^2)(q^{-1}+q^{-3})$	$\ominus \oplus \leadsto \emptyset$	$(1+q^{-2})(q+q^3)$

Identity: Has to do with the odd and even terms of Jacobi's triple product.

The identities we get have to do with *generalised Frobenius partitions* and *generalised Young diagrams*.

A nice three-state model:

Jump	Rate	Jump	Rate
$\oplus \rightarrow$	1	$\leftarrow \oplus$	q
$\leftarrow \ominus$	1	$\ominus \rightarrow$	q
$\emptyset \rightsquigarrow \ominus \oplus$	С	$\emptyset \leadsto \oplus \ominus$	qc
$\oplus \ominus \leadsto \emptyset$	2	$\ominus \oplus \leadsto \emptyset$	2q

Identity: Has to do with the square of Jacobi's triple product.

The identities we get have to do with *generalised Frobenius partitions* and *generalised Young diagrams*.

2-exclusion:

Jump	Rate	Jump	Rate
$\oplus \rightarrow$	1	$\leftarrow \oplus$	q
$\leftarrow \ominus$	1	$\ominus \rightarrow$	q
$\emptyset \leadsto \ominus \oplus$	1	$\emptyset \leadsto \oplus \ominus$	q
$\oplus \ominus \leadsto \emptyset$	1	$\ominus \oplus \leadsto \emptyset$	q

Identity: Looks new and interesting...

The identities we get have to do with *generalised Frobenius partitions* and *generalised Young diagrams*.

K-exclusion:

Identity: Rather nice generalisation using the Kth roots of unity.

The identities we get have to do with *generalised Frobenius partitions* and *generalised Young diagrams*.

K-exclusion:

Identity: Rather nice generalisation using the Kth roots of unity.

Zero range:

Identity: The geometric sum formula. :-D

The identities we get have to do with *generalised Frobenius partitions* and *generalised Young diagrams*.

K-exclusion:

Identity: Rather nice generalisation using the Kth roots of unity.

Zero range:

Identity: The geometric sum formula. :-D

Thank you.